

## Tactics and strategy of bioresonance and multiresonance therapy

A. Matison

(Firm "MATISONS" SIA, Riga, Latvia)

Term tactics is of Greek origin and originally denotes putting things in order. Word strategy also originates in Greek and is translated into Russian as the science of combat.

In order for our practice to be successful, it is necessary to pose and find answers to five questions:

- WHY?
- HOW?
- WHAT?
- WHERE?
- HOW?

The combination of honest and correct answers to the above questions allows the most correct application of the tactics and strategy of bioresonance (BRT) and multiresonance therapy (MRI).

Let's start with the first question: "WHY?". It would seem, at first glance, a rather stupid question. It must be placed in front of the patient. It may seem obvious why the patient came to you. But this is your opinion and your opinion. And you need to find out in detail and accurately from the patient, since, on the one hand, you find out the real task that the patient presents to you, and on the other hand, this serves as an oral agreement for the performance of work by you for which you receive a reward. On the part of the patient, these can be various tasks assigned to you, for example: he wants to find out the reasons for his ailment, find out the need to take this or that medicine, restore his own health. And in each specific case, you have to change tactics and strategy, based on the tasks that the patient sets for you.

Second question: "HOW?". It also has a degree of importance in that it allows you to most correctly assess the resources with which you will carry out the task assigned to you by the patient. The first and most important here are your knowledge, skill, experience and understanding, as well as the tools at your disposal. The best is the equipment produced by the IMEDIS Center - APK IMEDIS-EXTERT, MINIEKSPERT-D, a device for color therapy, the ASTROMED-M program. Knowledge of homeopathy and good programs for homeopathic repertorization of the patient are highly desirable as well.

Third question: "WHAT?". First of all, this is a question to yourself:

- what I need to understand in the patient,
- what motivates the patient to come to me,
- what is the source of the patient's problems,
- what the patient needs to understand in himself,
- what is important and useful for the patient that makes him a problem
- what needs to be changed in order to restore order, and bring the body to harmony and health.

All this is necessary to find out the reasons. To do this, it is best to use all diagnostic methods available to you, as classic

clinical and EPD methods, such as segmental diagnostics, which will allow you to assess the state of the autonomic nervous system, to some extent to even out the energy balance and then you can do without quadrant measurements, and at the same time optimize the indicators.

EPD by the method of R. Voll, in which it is best to start by measuring the chakra points. This allows you to estimate the level, or floor, on which the greatest deviation from the norm occurred. If you do not want to measure all points, then you will get the most optimal picture of the patient's condition by measuring the end and control points of the meridians. It is desirable to measure all points of the meridian with maximum deviations, especially if these deviations are below normal. At the same time, you need to understand that measurement by R. Voll's method is ascertaining and rather strongly subject to fluctuations that you can exert on the quality of measurement with your mental field.

The next stage of diagnostics is measurement according to the method ART or better yet - VRT +, at the same time, it should be understood that this method is the most accurate, since it is load-bearing and is less susceptible to outside influences, and allows not only to ascertain, but to identify the cause and primary source of the problems. If you have an apparatus for ART +, then you can immediately start measuring at the second level, which shows the state of the cell membrane. The first level of measurement on the apparatus for ART + correlates with the measurement according to the method of R. Voll, which mainly show the state of the intercellular space. Having measured the state of biological indices, the adaptation reserves at the second level, having identified the most affected and primarily affected organs, one can proceed to measuring at the third level, which shows the state of the cell cytoplasm. It is more correct to measure the indicators of cell metabolism at this level, since they are more in line with reality. Equally important at this level of measurement is the identification of the affected meridians, the identification of the primary affected meridian, the primary affected point on the primary affected meridian. At this level, it is possible to identify the viral infection of the body and the organ most affected by viruses. It must be remembered that it will be more correct to identify bacterial, mycotic and helminthic lesions at the second (first) level. Since these parasites live outside the cell, you need to look for them in the same place. Viruses live inside the cell, and therefore they need to be looked for at the third level of measurements, and since they can be incorporated into the structure of RNA and DNA, they can also be detected at the fourth level, which shows the state of the cell nucleus and its chromosomal apparatus. An important dimension at the fourth level is the identification of damaged chromosomes, and here you can also identify the agent that damages them. This may be the most common damage recently encountered by radiation, embedded viruses, viroids or their parts, and if they are not tested, then it is possible to identify the levorotatory amino acid (s) and find the primary significant one using a pointer to the primary damage (Zn met D400), which will restore the correct structure of the chromosome. A list of all these measurements allows you to determine at what level the greatest changes have occurred. Do they relate only to the information level, which determine changes in indicators of psychological stress, functional or organomorphological levels, or the changes have already affected everyone, or viroids or their parts, and if they are not tested, then it is possible to identify the levorotatory amino acid (s) and find the primary significant one using a pointer to the primary damage (Zn met D400), which will restore the correct structure of the chromosome. A list of all these measurements allows you to determine at what level the greatest changes have occurred. Do they relate only to the information level, which determine changes in indicators of psychological stress, functional or organomorphological levels, or the changes have already affected everyone, or viroids or their parts, and if they are not tested, then it is possible to identify the levorotatory amino acid (s) and find the primary significant one using a pointer to the primary damage (Zn met D400), which will restore the correct structure of the chromosome. A list of all these measurements allows you to determine at what level the greatest changes have occurred. Do they relate only to the information level, which determine changes in indicators of psychological stress, functional or organomorphological levels, or the changes have already affected everyone, or at what level the greatest changes have occurred. Do they relate only to the information level, which determine changes in indicators of psychological stress, functional or organomorphological levels, or the changes have already affected everyone, or at what level the greatest changes have occurred. Do they relate only to the information level, which determine changes in indicators of psychological stress, functional or organomorphological levels, or the changes have already affected everyone, or

several levels in various combinations. All measurements made allow a comprehensive assessment of the patient's condition and move on to the next question

"WHERE?" there is a lesion focus, which is the primary damaging aspect. This question is closely related to the question of selectivity, where our influence will be directed to achieve the goal set for us by the patient. Where to place the electrodes and where to bring the corrective impact (BRT own or external vibrations, MRI, This color therapy, acupuncture, homeopathy). brings us to the next question

"HOW?", when and how you will act on the cause of the problem. This, first of all, again depends on your equipment, what devices for the impact you have. If you have the very first self-contained apparatus for BRT, then you can choose the correct temporary exposure and the place of applying the electrodes on it. If you have a device of the second generation APK "BRT-VOLL", then the level of selectivity on it can be much higher. But with any treatment with BRT or MRI, it will be correct to exclude the meridian, which is in depletion or degeneration, from the exposure circuit. Choose a program for influencing an organ, system, therapy in the mode of brain rhythms or according to nosologies from the programs presented in the APC, depending on the problem being solved. In this case, it is useful to use the ASTROMED-M program to determine the strategy for recording a BR preparation.

to be guided by chronobiological rhythms and not to impose electromagnetic oscillations on the body, the relevance of which is no longer present. If you own the APK "IMEDIS-EXPERT", or the devices of the MINI-EXPERTDT series and the drug selector, which allow you to more selectively approach the correction of the problem, then you bear a greater degree of responsibility, since the effect on the patient during multiresonance therapy is more profound, and an error in its incorrect appointment can lead to more sad consequences, since we can transfer the problem from the visible for our diagnostic equipment to a deeper level, which we cannot recognize using measurements according to the method of R. Voll and even with the help of ART, which can lead us to a sense of well-being (imaginary), while the pathological process is gaining momentum, but at a deeper and more dangerous level for the patient's life - the level of impregnation of the cell cytoplasm, which is more difficult to diagnose and treat. In homotoxicology, this process is called progressive vicarization, and in homeopathy, anti-Hering's law. Our influence should be directed at the process of regressive vicarization, or the release of the cell, the intercellular space and the body as a whole from toxins and associated electromagnetic oscillations disharmonious for the body. APK "IMEDIS-EXPERT" allows Our influence should be directed at the process of regressive vicarization, or the release of the cell, the intercellular space and the body as a whole from toxins and associated electromagnetic oscillations disharmonious for the body. APK "IMEDIS-EXPERT" allows Our influence should be directed at the process of regressive vicarization, or the release of the cell, the intercellular space and the body as a whole from toxins and associated electromagnetic oscillations disharmonious for the body. APK "IMEDIS-EXPERT" allows perform BRT, MRI, electrotherapy (which is very often forgotten) and multiresonance therapy. The most successful is the use of multiresonant therapy using potentiated (homeopathic) medicines (electronic copies). It should be remembered that any type of BRT is aimed at suppressing pathological (disharmonious)

fluctuations, and this action can be regarded as an action in the plane, and the action of homeopathic remedies is directed inward, reaching the level of chromosomes. And like a scalpel revealing a psychoemotional problem the occurrence of pathology. The action of the potentiated drug has two phases of its action: 1) the emergence of a drug disease, 2) the mobilization of all the body's defenses aimed at eliminating the consequences of this drug disease and at the same time, according to the degree of similarity, eliminating the patient's problems. Choosing the right medication is essential in a patient's treatment strategy. APK "IMEDIS-VOLL" allows you to search for this only potentiated drug at a fundamentally different level, devoid of the influence of subjectivity both on the part of the patient and the doctor.

If you have a patient who is ill for a long time, then this may indicate that there is a stagnation of both Chi energy and hemodynamics in his body. This, in turn, can lead to adhesion of blood corpuscles, which reduces their useful sorption surface and leads to metabolic disorders, which can manifest itself in the form of pastiness, edema, inflammation, degeneration, which we can determine using the ART + method (ART ). In this case, it is more correct to use MRI through various inductors, depending on the size and area of the problem. MRI is also convenient to apply to young children, for whom the problem is to sit quietly for at least a few minutes. MRI is also applicable in veterinary practice. MRI can be combined with a properly selected potentiated medication. If you work in the Windows environment, then you can immediately combine into MRI, BRT and homeopathic action. But remember that more is not necessarily better. And the level of your professionalism (understanding) allows you to achieve maximum therapeutic effect with minimal exposure. This is the art of healing. With any magnetic effect, an electromagnetic wave is generated, which contributes to

normalization of the electrical potential of the cell membrane, disintegration of cells and an increase in the contacting (working) surface of the cell, increased blood flow rate, opening of capillary collaterals. This, in turn, normalizes homeostasis.

Electrotherapy is indicated in the same cases where conventional physiotherapy is prescribed. With the correct application of electrodes, it can contribute to myostimulation or stimulation of nerve receptors, the restoration of the work of both striated and smooth muscles, and you can also use the method of stimulating points, which was proposed by R. Voll.

Important for any method of treatment is the normalization of indicators at points and, especially, on chakra measurements by the method of R. Voll, as well as an increase in the indicators of adaptation resources, and bringing the indicators of biological indices to optimal at all measurement levels by the ART + method.

It is also necessary to pay attention to the use of color therapy. It is important that a healthy body needs light, and a sick one needs color. The transfer of information in our body is carried out with the help of light quanta, along the meridian channels, which are structures of our body superconducting for light quanta, through which the transfer is carried out

information from cell to cell. All biochemical processes in the cell are accompanied by photonic activity. Therefore, the most important application of color therapy is pathology associated with the third (cytoplasmic) level of damage to the body. Correctly selected color and its selective effect on the initially damaged acupuncture point, with the help of a color therapy apparatus, allow eliminating the indicators of the pathological process revealed during the examination in a matter of seconds. With this method of exposure, it is very easy to overshoot the optimal level of exposure, and therefore it must be carefully controlled using VRT (+), using the Cu met D400 optimization index.

How often and for how long should BRT, MRI and multiresonance therapy be performed? It depends on situation. In each case, you need to be approached individually. It can be both short-term courses and a single exposure. In severe conditions, continuously until recovery normalization of vital functions. Everything is dictated by the circumstances. If color point light therapy is used, this is a few seconds. The use of BRT and MRI per point takes 10–20 minutes. The use of BRT and MRI for general strategies or brain rhythms - from 20 min. and more. In this case, one should be guided by the fact that the synthesis of protein molecules requires at least 20 minutes. and more.

It is possible to prolong the normalizing effect of therapy by prescribing a BR drug or by prescribing a potentiated (homeopathic) medication. Moreover, if the potency and amount of the drug can be determined using EPD, then the frequency, duration and time of admission using the method of R. Voll and ART is difficult to determine. To do this, you can use such a type of diagnosis as mental testing, the use of a biotensor, or a muscle test.

The quality of your impact will be determined by the accuracy of finding the nucleus of the pathology, determining the underlying factor that gave rise to this nucleus, determining the drug that can solve this problem, as well as the method (s) of exposure and their combination. It can be a point, segment, area, or the whole organism. And try not to forget that the smaller and more precise the degree of exposure, the better the therapeutic effect.

The most important thing is the condition of the patient in front of us, and not the indicators of the instruments.

Control the methods of diagnosis and exposure used by you, clinical methods of research of the patient. Each repeated appointment should be started with a detailed examination of the patient, maybe in a slightly more concise version, and it is advisable to carry out diagnostics using the ART + method. This will allow, indeed, to control the healing process, and not gloss over the symptoms, which is so often the case with modern medicine.

It is necessary to help the patient to heal, that is, to cure himself, and the doctor in this case can only help, and not do it for the patient.