

Use of targeted blood nosode (NANCr) and systemic spiritual adapters as a way to optimize the healing process

D.B. Okun, I.B. Okun, K.N. Mkhitarian
(LPC "Synergy", Vladivostok; Center "IMEDIS", Moscow,
Russia)

In clinical practice, situations arise when a specialist needs to provide full assistance to a patient in a fairly short period of time, i.e. "Prepare" a drug that would completely stop the main pathological process and, at the same time, would not remove the entire system from the state of adaptive equilibrium.

As a research model, the methodological developments of A.E. Kudaev, K.N. Mkhitarian, N.K. Khodareva were used. on the preparation of a targeted blood nosode and the use of systemic spiritual adapters (NANCr + SDA).

For 6 months, all patients (86 people) who sought medical help underwent therapy in the NANCr + SDA mode. As a target marker, the sum was used: organ-specific (OSP) drugs and nosodes of both infectious and non-infectious series, and others that have a resonant relationship with the OSP and nosode (disease). The choice of SDA was carried out "directly" followed by "recording" and prescription as a separate drug.

In the course of therapy, all patients showed positive dynamics: an improvement in the quality of life, an absence or a significant decrease in intensity of manifestation of the main complaints, stabilization of the emotional background and normalization of laboratory parameters. In each case, the frequency and single dosage were selected individually. The average frequency of admission (if necessary) was 25–35 days.

Clinical examples

1. Patient F., 20 years old, neuritis of the facial nerve, exacerbation. Within 4 days there is a virtual lack of effect from the use of drug therapy. After taking NANCr and SDA, an exacerbation occurred - an increase in pain syndrome within 2 minutes, followed by a sharp improvement. After the second dose of the drug, the exacerbation was repeated, but with a lower intensity of the process, after which there was a significant improvement in the condition with a completely arrested pain syndrome and the absence of neurological abnormalities.

2. Patient V., 65 years old, incipient cataract. Against the background of application NANCr and SDA have a stable positive effect - the results of the examination by an ophthalmologist.

3. Patient I., 44 years old, bronchial asthma, mixed form, the number of seizures up to 18 times a day, after taking NANCr and SDA, the number and severity of seizures decreased, and on the 25th day seizures were completely absent.

4. Patient A., 44 years old, hypertension, crisis course. On the against the background of this therapy, there is a persistent decrease in blood pressure of the order of 5–8 mm. rt. Art. per month, crises were not observed, the number of medications was reduced from 4 to 1, observation and therapy continues.

5. Patient 38 years old, exacerbation of chronic orcho-epididymitis, appointed NANCr and SDA - admission within 20 days, the conclusion of the urologist - healthy.

Conclusion: the use of NANCr and SDA is a fairly effective method for a wide variety of pathologies with a stable positive effect. I would like to note that the use of this technique allows to compensate not only the manifestations of the underlying disease, but also has a systemic effect - the normalization of hemodynamic parameters, sleep, appetite and psycho-emotional sphere.

D.B. Okun, I.B. Okun, K.N. Mkhitarian Using targeted blood nosode (NANCr) and systemic spiritual adaptants as a way to optimize the healing process // XIII