

Case from practice

S.Yu. Kuznetsova

(LLC "Center for Energy Information Medicine", Samara, Russia)

Patient A., 13 years old, turned to CEIM on 15.02.2005 with complaints of attacks of loss of consciousness, accompanied by convulsions, biting the tongue, blanching of the skin, lasting 2-3 minutes, preceded by cramping pains in the umbilical region. Periodically attacks of loss of consciousness are accompanied by involuntary urination. Such attacks occur 1 time in 2 months from the age of three. The patient is periodically disturbed by pain in the left hypochondrium after eating fatty foods. In addition, the patient has acne on the face and back.

History of frequent acute respiratory viral infections. At 7 months - a concussion. At the age of 3, an attack of loss of consciousness first occurred. Examined by a neurologist, performed an EEG, diagnosed with temporal lobe epilepsy. In this regard, he is registered with a neurologist, constantly takes antiepileptic drugs. However, the treatment does not bring a special effect; against the background of conservative therapy, epileptic seizures continue to occur.

Diagnostics using the ART method revealed:

- geopathogenic load 3 tbsp.,
- mental load of 7 tbsp.,
- the average degree of depletion of the immune system,
- bacterial and fungal intestinal dysbiosis,
- bacterial (pneumococci D8, streptococci D5) and fungal burden in the bronchi, palatine tonsils,
- fungal burden in the pancreas,
- intestinal giardiasis D15,
- trichinosis D4, ascariasis,
- toxoplasmosis D32.

The child is prescribed therapy:

- 1) Therapeutic diet.
- 2) Treatment with private and general BR-drugs (through the optimal step on adaptation reserves).
- 3) Therapy fixed frequencies (antiparasitic, antifungal and antibacterial), incl. 6, 2 Hz.
- 4) Treatment with nosodes in conjunction with a complex of drainage preparations firm "ONOM".
- 5) Preparations "Bach Flowers".
- 6) Hilak-forte and bifidumbacterin.
- 7) Sanitation of the lacunae of the palatine tonsils.

After the treatment, over the next two years, there were no epileptic seizures, no acne, no pain in the abdomen. Does not take antiepileptic drugs.