

Statistical data obtained when working by the method of vegetative resonance test
 "IMEDIS-TEST +" (results of work from May 2003 to February 2007)
 G.Yu. Andreichenko
 (Vorkuta, Russia)

In total, during this period, a four-level diagnosis of 828 people was carried out. By age, the groups were distributed as follows.

Table 1

Age	Children under 1 of the year	1-10 years	11-20 years old	21-40 years old	41-60 years old	Over 60 years old	Total
Fact. qty	eleven	121	115	241	301	39	828
%	1.3%	14.6%	13.9%	29.1%	36.4%	4.7%	100%

The current picture of the dominant miasms is presented in Table 2

table 2

Dominant miasm	Fact. number of people	%
Botulism	7	0.9
Toxoplasma	17	2.0
Borrelia	4	0.5
Tuberculosis	35	4.2
Rickettsia	one	0.1
anthrax	one	0.1
Fungoid fungus	49	5.9
Coxsackie virus	6	0.7
Epstein-Barr virus	five	0.6
Cytomegalovirus	119	14.4
Hepatitis	76	9.2
Herpes	12	1.5
Rabies	3	0,4
Measles	10	1,2
Smallpox natural	10	1,2
Chicken pox	35	4.2
Polio	252	30.5
Rubella	141	17.0
Mycoplasma	2	0.2
Diphtheria	3	0,4
Syphilis	one	0.1
Pneumocystus	one	0.1
Plasmodium malaria	one	0.1
Flu	one	0.1
Psychovegetary load	36	4.4
TOTAL	828	100.0

The first 7 places in descending order are occupied by the dominant miasms of poliomyelitis viruses (30.5%), rubella (17.0%), cytomegalovirus (14.4%), hepatitis (9.2%), fungoid fungus (5.9%) , chickenpox (4.2%), tuberculosis (4.2%). That is 85.4% of the total.

In 36 cases, at the fourth level of ART +, only psycho-vegetative loads were tested, which were eliminated with drugs from the Bach Flowers group, Systemic Spiritual Adaptants, and MEDFARMA (psychosocial loads).

In 137 people, at the fourth level, miasm and psycho-vegetative loads were determined, which were removed by Viper 2000 in combination with a drug from the Bach Flowers group. Moreover, the drug from the Bach Flowers group served in this case as a drainage.

Differentiating through the Intox II and III pointers, the following picture emerged (see Table 3).

Table 3

Miasm	Intox II	Intox III	Total
Botulism	-	7	7
Toxoplasma	one	sixteen	17
Borrelia	one	3	4
Tuberculosis	-	35	35
Rickettsia	-	one	one
anthrax	-	one	one
Fungoid fungus	-	49	49
Coxsackie virus	-	6	6
Epstein-Barra	-	five	five
Cytomegalovirus	3	116	119
Hepatitis	4	72	76
Herpes	one	eleven	12
Rabies	-	3	3
Measles	one	nine	10
Smallpox natural	-	10	10
Chicken pox	five	thirty	35
Polio	five	247	252
Rubella	6	135	141
Mycoplasma	-	2	2
Diphtheria	-	3	3
Syphilis	-	one	one
Pneumocystus	-	one	one
Plasmodium painter.	-	one	one
Flu	-	one	one
TOTAL	27-3.4%	765-96.6%	792-100%

When analyzing the diagnostic picture, there is a clear predominance of hereditary burden (96.6%).

For the therapy of level 4 ART +, I used the following drugs:

- special Viper 2000 - 775 cases (93.6%),
- FM chords - 16 cases (1.9%),
- nosode - 1 case (0.1%),
- "Bach Flowers" - 25 cases (3.0%),
- Systemic Spiritual Adapters - 7 cases (0.9%),
- other drugs - 4 cases (0.5%).

Over the past year, there has been a 2.35-fold increase in the number of patients in whom only psycho-vegetative loads are determined at the fourth diagnostic level of ART +. Solving these problems with drugs from the Bach Flowers, Systemic Spiritual Adaptants, Medpharma it is possible to remove the autoaggressive program in DNA (see table 4).

Table 4

Year	2004 r.	2005 year	2006 year	2007 year
Fact. qty	nine	10	13	36
%	2.4%	1.8%	1.9%	4.4%

After removing the miasms, I conduct therapy for deep mycoses.

A few figures on the results of antifungal therapy. In total, 491 people passed this stage. Directly 607 sessions were carried out. The most frequently tested fungi are listed in Table 5.

Table 5

Type of fungus	First session	Repeated session	Total	%
Fungoid fungus	185	68	253	41,7
Dermatomycosis	62	6	68	11.2
Aspergillosis	35	4	39	6.4
Hansenulla	13	-	13	2.1
Rodoturool	10	2	12	2.0
Pihia pini	13	2	fifteen	2.5
Saccharomycetes	12	one	13	2.1
Mukor pusilus	eight	-	eight	1,3
Schwaniomycetes	4	-	4	0.7
ctinomycosis	4	-	4	0.7
Candidiasis	2	-	2	0.3
Other (meetings. 1-3 times)	143	33	176	29.0
TOTAL	491	116	607	100%

G.Yu. Andreichenko Statistical data obtained by the method of vegetative resonance test "IMEDIS-TEST +" (results of work from May 2003 to February 2007) // XIII "- M .:" IMEDIS ", 2007, vol.2 - p.216-220