

The use of induction therapy in patients with psychiatric pathology

T.V. Dondyk

(Moscow Homeopathic Center, Moscow, Russia)

Practical experience in treating patients under stress, with behavioral and emotional disorders, manifestation of schizophrenia, MDP, and depressive disorders has shown high efficacy and safety of induction therapy and multiresonance therapy compared to allopathic treatment. In the course of therapy, frequencies corresponding to the rhythms of the brain were used - alpha, beta, theta, delta in the swing mode and frequencies of 40 Hz, 38 Hz. The frequencies required for the treatment were preliminarily tested by the ART method using the IMEDIS equipment. The intervals between therapy sessions are from 1 to 5 days. Patients were prescribed BR drugs after BR therapy according to strategies 2, 3, 4.

The choice of frequencies 38 and 40 Hz was based on the studies of Ivanitskiy A.M., in particular, on the method he developed for mapping intracortical connections. It was found that intracortical connections are carried out at a frequency of exactly 40 Hz. In cognitive tests, intercortical interaction is carried out at a frequency of 38 Hz.

A number of authors attach great importance to the synchronization of potentials at a frequency of 40 Hz in the mechanisms of consciousness and the formation of visual images. In pathology, the interaction between different parts of the cerebral cortex is disrupted. So, schizophrenia is recognized as a disease associated mainly not with damage to certain brain structures, but with a violation of the relationship between these structures, with their functional disintegration. When performing a cognitive task, the number of interhemispheric connections in healthy people increases to eight. In patients with schizophrenia, using the method of mapping intracortical connections, it was possible to reveal the complete absence of interhemispheric connections. In acute schizophrenia (with positive symptoms), there is a violation of the system of interhemispheric connections, which can still be restored, and with proper treatment, hope for a good prognosis. In chronic schizophrenia (with negative symptoms), a new pathological network of interhemispheric connections at a low frequency already arises. It is more difficult to destroy the network and the forecast in this case is much worse. In depression, there is an asymmetry in the indices of intercortical connections.

ART showed resonance of frequencies corresponding to theta, delta, alpha and beta rhythms, 40, 38 Hz in patients with epilepsy, schizophrenia, depression, under stress conditions. Taking into account the positive dynamics in treatment, it can be assumed that in the process of induction therapy with the help of a magnetic inductor - a device for magnetic therapy from the firm "IMEDIS", there is a synchronization of intercortical interactions and an increase in their number. This method allows the treatment of patients with neurological and mental pathology without the use of allopathic drugs that are detrimental to

body side effects. In the treatment of patients who use psychotropic drugs for a long time, BRT allows to reduce the doses and side effects of allopathic drugs.

Clinical examples

1. Irina Z., 27 y. Diagnosis: Manic-depressive psychosis.

The duration of the disease is 1.5 years. Aggravation. Within 4 months she took Rispolept 4.5 mg, Akinetone 2.5 mg. At the initial admission, it is inadequate, inhibited, fixed on the disease, during testing, a pronounced degree of depression, high anxiety are determined. There is stiffness of muscles, absence of menstruation (within 3 months). Induction therapy was carried out, frequencies of 40 and 38 Hz were used (sessions 2 times a week), BR-drugs were prescribed. After 2 months. the patient's psychosomatic status returned to normal, the drugs were canceled (the drugs were canceled gradually). Follow-up for 5 years. I. is still adequate and successfully found a job.

2. K., 68 years old. Complaints of frightening auditory hallucinations, fear leave the apartment. Persecution delirium. The duration of the disease is 2 months. The patient's daughter decided not to go to psychiatrists, but to try more gentle treatment. After 15 sessions of therapy, 2 times a week, the mental state returned to normal. Follow-up - 7 years.

3. M., 9 years old. The boy is sluggish, has difficulty reading, at school, the reading score is "2". The level of impaired psychological adaptation according to the Luscher test is 113. After 12 sessions of therapy, the child became more active, communicative (this was noted by both mother and teachers at school, and other people close to the family), the reading speed increased from 18 to 38 words per minute. The level of impairment of psychological adaptation according to the Luscher test decreased to 83. Repeated tests "Drawing of a person", "Drawing of a family" with obvious positive dynamics.

4. K., 27 years old. After hypnogenic influence - delirium of persecution, attempted suicide, disorientation in the outside world. One session of induction therapy and subsequent administration of bioresonance drugs for a week normalized the condition. Follow-up - 7 years.

5.D., 10 years old. Complaints of twitching of the nose and left hand. Carried out 6 sessions of induction therapy and taking bioresonance drugs for a week normalized the condition. Follow-up - 7 years.

6.S., 44 g. Diagnosis: endogenous depression, was unsuccessfully treated in within 6 months. in a psychiatric clinic. Therapy using induction programs, 38 Hz against the background of psychotherapy, taking bioresonance drugs for 1 month completely normalized the mental state. S. got a job. Follow-up - 4 years.

7. When adding new fish to the aquarium, two of the old ones got sick.

Outwardly, the disease was regarded as a fungal infection (cloudy eyes, full of holes, covered with white fins). After the death of one fish, 1 aluminum glass of water was added to the aquarium with the frequency recorded on it on the device of the IMEDIS Center of the anti-fungal 1 and 2 drugs. Therapy lasted 10 days with the addition of the drug to the water once every 23 days. As a result, the fins of the diseased fish completely recovered, the cloudy spot on the eye decreased by 2 times. Follow-up - 5 months

Assuming the absence of a placebo effect on the part of the patient, it can be argued that energy-informational copies of drugs work even without a psychotherapeutic component.

These examples show that this type of therapy makes it possible to carry out symptomatic and etiopathogenetic treatment in patients with psychiatric pathology, sometimes more successfully and safely for the patient in comparison with allopathic treatment. This direction needs to be developed and studied. However, we must not forget about the need for subsequent psychological rehabilitation by methods of individual, family psychotherapy, homeopathic preparations, allowing people who get rid of symptoms to look at the world around them in a different way.

Literature

1. Alexandrov Yu.A. Psychophysiology. - 2001
2. Samokhin A.V., Gotovsky Yu.V. Electropuncture diagnostics and Voll therapy. - M.: IMEDIS, 1995.
3. Gotovsky Yu.V., Kosareva LB, Frolova LA. Quick Start Guide on induction therapy. - M.: IMEDIS.
4. Savchenko A.B. Central and peripheral effects transcranial electrical stimulation. - 1994.
5. Lomarev M.P. Structural and functional rearrangements in the human brain during therapeutic transcranial electrical influences with pulsed and direct current. - 1995.
6. Pakhomova Zh.M., Bogdanov O.V. Influence of transcranial micropolarization of the frontal cortex on the state of motor and cognitive functions in extrapyramidal pathology.

T.V. Dondik Application of induction therapy in patients with psychiatric pathology // XIII