Bioresonance therapy for headaches of vascular origin A.V. Shamanin (LPTs "Sinegria", Vladivostok, Russia)

Headaches of vascular origin are types of headaches that are quite common in practice. Dysfunction of the vascular component is, in most cases, secondary.

In the ART system, specific tests for the diagnosis of this pathology have not yet been developed, and the use of tests for the presence of toxins and free radicals and vegetative burden on the vascular component of the brain is not always adequate, because:

- the problem is often of a functional nature;
- patients come to the appointment without active complaints.

The test "toxins and free radicals" is tested through an organopreparation (OSP) of arteries in low potencies, and they do not reflect vascular spasm, but rather reflect paralytic processes in the arteries (vascular spasm as such is associated in the ART system with organopreparation of arteries of high potencies).

Analysis of 86 case histories revealed the following pattern: it is most appropriate to use indicators of autonomic dysfunction (tension or depletion of the sympathetic part of the ANS) and look for their resonance with the vascular component of the brain.

In my practice, I use organopreparations in single potencies - the arterial network of the brain, paravertebral arteries, the pia mater. The potencies revealed on these organs, corresponding to the dysfunctions of the sympathetic division of the ANS, are recorded on homeopathic globules and placed outside the contour, as a single pathophysiological index (PU), then all factors can be viewed on the ART test (external loads, infectious nosodes, deficits, etc.) etc., which are used to prepare the second preparation). After that, we test through the PU filter all organ products of the brain and the periphery that resonate with this indicator - these are the organs that are combined into a single pathophysiological circle, some of them are the organs of discharge, some are victims, and some are those that try this situation if not to correct,

we get OSBres. It is our task to fix or destroy this circle.

After that, everything found (PU + Σ OSP_{res}) we carry out on the homeopathic crumbs for the contour and carry out BRT with frontal leads at the same time mode along all meridians, "golden section", the patient undergoes therapy until the response, after which we write down the BR-drug "main". The second drug is prepared according to generally accepted methods for previously identified loads.

This technique has shown its high efficiency in a larger number of patients, up to 75%, during 3-5 sessions with a frequency of 1 time per month, it is possible to either reduce headaches to a minimum or completely get rid of them.

Work on optimizing the therapeutic approach for this pathological condition continues.

A.V. Shamanin Bioresonance therapy of headaches of vascular origin // XIII

Imedis conferance 2007 NO: 265		