

## Male infertility: problems of diagnosis and treatment

Popova E.Yu.

("Center for Energy Information Medicine", Samara, Russia)

Infertility is one of the most pressing and difficult medical and psychological problems ... Russian statistics are sad - every fifth couple of childbearing age is infertile today. But even these numbers are possibly inaccurate. The difficulties of counting are connected with the fact that people learn about their infertility only when they have a desire to have children, and this "does not work." Many people with impaired fertility are simply unaware of their illness, since it does not affect their vital interests.

Infertility, as a medical diagnosis, is said when a couple of childbearing age who does not use contraception do not become pregnant after one year of regular sexual activity.

To date, many factors have been identified that lead to a violation of human reproductive capacity. Women have them 22, in men - 16. Often the infertility of a couple is caused by complex factors (male and female infertility), according to the international classification of diseases, they are considered as two different diseases.

The main diseases that lead to infertility in women are inflammation of the uterus and its appendages, dysfunction of the ovaries, endometriosis, and many others.

We were interested in the problem of male infertility, tk. In almost 50% of cases, when examining a married couple, it turns out that the husband suffers from some kind of chronic urological disease and his fertility (ability to fertilize) is significantly reduced, and sometimes reduced to zero, therefore, in parallel with his wife, the man should be examined, and if it is necessary to undergo a course of treatment.

Male infertility is a disease in which a mature man is unable to reproduce offspring. WHO statistics from year to year provide disappointing information that in 45% of cases of infertile couples, the "culprit" is a man, and in 40% - a woman, the remaining cases are related

with incompatibility of partners (female body that suppress sperm activity) diseases. develops antibodies, and others rare

Modern medicine makes it possible to say with confidence that male infertility is a disease that must be treated.

The most common causes of male infertility are:

- inflammatory diseases of the genitourinary organs (urethritis, prostatitis and etc.);
- sexually transmitted diseases (gonorrhoea, chlamydia, trichomoniasis, etc.);
- sexual dysfunctions (impotence, premature retrograde ejaculation, etc.);
- obstruction (obstruction) of the vas deferens;
- expansion of the veins of the spermatic cord (varicocele);
- hormonal disorders leading to a decrease in spermatogenesis.

The Center for Energy Information Medicine has been operating in Samara for three years. We have developed and successfully operate a program for the diagnosis and treatment of infertility using the equipment of the firm "IMEDIS". For diagnostics, R. Voll's method and vegetative resonance test (ART) are used. Full diagnostics of both spouses, who applied to the Center, is carried out. It includes:

- examination of all systems, identification of foci of chronic infections;
- identification of electromagnetic, radioactive, geopathogenic loads;
- study of the endocrine system, hormone levels;
- identification of chronic diseases (especially of the genitourinary system) and their reasons;
- identification of the causes of infertility by the ART method;
- differential diagnosis of the type of male infertility (secretory, excretory, combined, immune, relative).

Infertility treatment is a long and rather complicated process and, according to our data, takes from 8 months. up to 1.5 years. The treatment is carried out in a comprehensive manner. Based on the diagnostics performed, an individual therapy plan is developed for each patient. The main methods of treatment are resonance frequency therapy and endogenous bioresonance therapy with the recording of private and general BR drugs.

Stage 1 - rehabilitation of foci of chronic infection, treatment of chronic diseases, restoration of hormonal levels and homeostasis.

Stage 2 - stimulation of spermatogenesis. For this purpose, in addition to bioresonance therapy, electronic analogs of homeopathic medicines and hormones are used (taking into account the cyclical nature of their formation in the body), selected individually by testing using ART.

At stage 2, massage of the prostate gland, acupuncture and electropuncture must be included. This is especially important with retrograde and premature ejaculation, which are often the result of autonomic dysfunction, osteochondrosis of the lumbosacral spine.

department

In about 10% of cases in the diagnosis of infertility in marriage medics come to a "dead end" - the spouses do not have any diseases that could interfere with fertilization. In this case, the equipment and methods of the "IMEDIS" company provide invaluable assistance to the doctor. As an illustration, I give a description of one case from practice.

In the fall of 2002, a married couple contacted CEIM about infertility. They have been married for 15 years, during which no pregnancy has occurred. The marriage is the first for both spouses. They were repeatedly examined in different medical centers, but to no avail.

Both spouses underwent complete electropunctural diagnostics according to the methods of R. Voll and vegetative resonance test. The wife was diagnosed with intestinal dysbiosis. No other obvious pathology was identified.

A careful study of her husband revealed: a decrease in the level of testosterone and progesterone, orchiepididymitis associated with maximum

DNA damage caused by the mumps virus. The patient categorically denies the disease with mumps. No other infections and diseases of the genitourinary system have been identified. Subsequently, with a careful collection of anamnesis, it was revealed that shortly before the patient's conception, his father suffered a severe form of mumps with testis lesions, which probably resulted in a distortion of DNA-related information.

In the spermatogram - a significant decrease in motile forms spermatozoa, a large number of immobile and sedentary forms, IgG (autoimmune process), an increase in the number of leukocytes. That is, the process of sperm maturation is disrupted.

- Within 1.5 years, several courses of therapy were carried out, including
- "erasure" of distorted hereditary information. For this purpose, we used electronic analogs of miasms from the company "OHOM", selected by testing;
  - endogenous bioresonance therapy with recording of private and general BRs; drugs;
  - resonant frequency therapy (anti-virus, normalizing hormonal background);
  - in order to stimulate spermatogenesis, electronic analogues of homeopathic drugs, hormones were prescribed (taking into account the cyclicity of their entry into the bloodstream in the body);
  - induction therapy;
  - electropuncture therapy.

As a result, the number of motile spermatozoa in the spermatogram increased to the lower limit of the norm, the signs of an autoimmune process disappeared.

A healthy boy was born to this couple on February 14.

Our Center has been operating in Samara for the fourth year. During this period, this is the fifth child born after the treatment of couples who came to us with the problem of infertility.

Thus, the equipment and progressive diagnostic methods of treatment offered by the IMEDIS Center give many married couples a chance to find the happiness of having children.