

Choosing a chiroglyphic palm line for performing HST
Gotovsky Yu.V., Mkhitaryan K.N.
(Center "IMEDIS", Moscow, Russia)

As a group MBAT, from which the body's response to the load is written off with a target marker (MC), usually one of the chiroglyphic lines of the palm is used. In this case, we say that: "XST is carried out according to corresponding chiroglyphic line ", which is a thus becomes chronosemantic line.

The criteria for choosing a chronosemantic lines for chronosemantic therapy (CHT) are the test results (for BAP measurements) of special chiroglyphic line selection markers. Each chiroglyph line corresponds to some marker of choice. The criterion for determining the chronosemantic line preferable for the XST using the selection marker is the following:

the reproducibility of the BAP on which the measurements are carried out.

The following can be used as sets of chiroglyphic line selection markers:

1. A set of seven alchemical (known to alchemists) metals in thousandth potencies:

Aurum met. C1000 (gold) - marker for choosing the line of the Sun (Apollo);

Argentummet. C1000 (silver) - marker for selecting the lines of the Moon / Uranus;

Mercurius met. C1000 (mercury) - marker of the choice of the line of Mercury (Liver, Health);

Cuprummet. C1000 (copper) - marker for choosing the line of Venus (Life);

Ferrum met. C1000 (iron) - marker for selecting the line of Mars (Head);

Stannummet. C1000 (tin) - marker for choosing the line of Jupiter (Heart);

Plumbum. met. C1000 (lead) - marker of the choice of the line of Saturn (Destiny).

2. A set of markers for the preference of one or another planetary functional, used by the center "Synergy" (Vladivostok):

symbolic script marker (MSS) "Chronosemantic functional of the Sun" 1,2 marker for choosing the line of the Sun (Apollo);

MSS "Chronosemantic functional of the Moon" marker of the choice of the line of the Moon (Travel) / Uranus;

MSS "Chronosemantic functional of Mercury" 1.2 marker of the choice of the line of Mercury (Liver, Health);

MSS "Chronosemantic functional of Venus" 1,2,3 marker of the choice of the line of Venus (Life);

MSS "Chronosemantic functional of Mars" 1.2 marker of selection of the line of Mars (Head);

MSS "Chronosemantic functional of Jupiter" 1.2 Jupiter (Heart) line selection marker;

MSS "Chronosemantic functional of Saturn" 1.2 marker of the choice of the line of Saturn (Destiny);

MSS "Chronosemantic functional of Uranus" marker of the choice of the line of the Moon / Uranus;

MSS "Chronosemantic functional of Neptune" 1.2 Neptune line selection marker;

MSS "Chronosemantic functional of Pluto" 1,2,3 Pluto line selection marker.

The final choice of the chiroglyphic line, along which the CTT is performed, remains, of course, with the doctor. An alternative strategy for choosing a chiroglyphic line for CT is to carry out CT along all major chiroglyphic lines (OHL) of the palm or on OHL belonging to the horizontal or vertical bundle of chiroglyph lines. In this case, it is usually advisable to conduct therapy in the following sequence: chiroglyphic lines of Venus (Life), Mars (Head), Jupiter (Heart), Saturn (Destiny), Sun (Apollo), Mercury (Liver, Health), Moon (Travel) / Uranus.

3. A set of electromagnetic oscillations written off from the OHL endpoints:

VenD_{max} marker of the choice of the line of Venus (Life);

SatD_{max} marker of the choice of the line of Saturn (Destiny);

SanD_{max} marker for choosing the line of the Sun (Apollo);

MerD_{max} marker of the choice of the line of Mercury (Liver, Health);

MarD_{max} marker of selection of the line of Mars (Head);

JupD_{max} Jupiter (Heart) line selection marker;

LunD_{max} marker of the choice of the line of the Moon (Travel) / Uranus.

Each of the developed sets of markers for choosing one of the OHL has your "area of preference".

Set 1 is used to select OHL in cases where mental symptoms are the leading symptoms by which the nosology and the preferred route of treatment are classified.

Set 2 is used to select OHL in cases where somatic symptoms are the leading symptoms according to which nosology and the preferred route of treatment are classified.

Set 3 is used to select OHL in cases where the leading symptoms according to which the nosology and the preferred way of treatment are classified are mixed symptoms that cannot be attributed to either mental or nosological.

In the volumetric course of the XST, it is advisable to alternate the markers of the choice of OHL.

Gotovsky Yu.V., Mkhitaryan K.N. Choosing a chiroglyphic palm line for the implementation of XST // X

:" IMEDIS ", 2004, vol.2 - p.214-217