

Diagnosis and treatment of endometriosis according to the method of Professor A.A. Hovsepyan using the apparatus "MINI-EXPERT-D"
Selyutina N.I., Bessarabov O.V.
(Treatment and Diagnostic Center "Medicine of the Future", Kaliningrad, Russia)

Application of the methodology of Professor A.A. Hovsepyan in case of gynecological diseases, it allows to quickly and effectively identify the causes and eliminate the exacerbation of chronic diseases, in particular: endometriosis, endometritis and other inflammatory processes in gynecology. After the preparation of two or three BR-preparations, complaints of pain disappear, bleeding is eliminated, and the menstrual cycle is restored.

One of the main positive aspects of this therapy is that many women who have suffered from infertility for many years, after being treated in our center according to the method of Professor A.A. Hovsepyan. were able to get pregnant safely.

Example

Patient K., born in 1984. She complained of pain in the lower abdomen and persistent bleeding from the vagina. Pain, discharge, irregularities in the cycle appeared after an abortion in August 2004.

According to the ART + method and the method of Professor A.A. Hovsepyan revealed the presence of interference fields, indicating an acidic state, as well as a toxic focus in the uterus. According to the fluctuations taken from the toxic focus, the non-optimal state of the myometrium was revealed according to the following parameters:

- Myometrium,
- Anabolism 6 tbsp.,
- Acidity 6 tbsp.,
- Depletion of VNS 5 tbsp.,
- Sympathicus,
- Connective tissue insufficiency 2 tbsp.,
- Autoimmune processes 1 tbsp.

After testing Intox I, Intox II, Intox III, an acquired bacterial and mycotic load was revealed: chlamydia, Trichomonas, gonococci, ureaplasma, mycoplasma, candidiasis.

According to the degree of slugging of the mesenchyme, the most pathogenic pathogens were determined.

At the first session of bioresonance therapy, a therapeutic chain was established, aimed at correcting the ANS and metabolic processes. After the correction, resonance-frequency therapy was started for the most pathogenic pathogens, and the appropriate drains were selected. The dosage of the BR-drug was tested through the adaptation reserves.

In total, four sessions of bioresonance therapy with the preparation of four BR-preparations were needed to correct the ANS, establish the necessary metabolic processes, acid-base balance.

At the last session, BR-therapy was carried out to eliminate connective tissue insufficiency and autoimmune processes, a BR-drug was prepared,

which the patient took for 3 months.

After treatment, the condition stopped bleeding from the vagina, the menstrual cycle disappeared.

the patient returned to normal:
the pain went, returned to normal

Selyutina N.I., Bessarabov O.V. Diagnosis and treatment of endometriosis according to the method of Professor A.A. Hovsepyan using the device "MINI-EXPERT-D" // XI "IMEDIS ", 2005, v.1 - C.212-213