## Clinical examples of curing cancer patients by methods information medicine Gorbenko S.V. (West Kazakhstan Regional Oncological Dispensary, Uralsk, Kazakhstan)

## 1. LVB, born in 1964, map number 1215. Date of diagnosis 06.06.98. Diagnosis: myogenic sarcoma of the right hip stage IV. X-ray from 07/13/98: multiple metastases to the lungs. Cytology of tumor punctate from 06.06.98: myogenic sarcoma.

Due to the obvious futility of standard chemotherapy, the patient was offered parallel treatment with bioresonance therapy. The diameter of the femoral tumor was measured before the start of treatment. Within 10 seconds, information from the tumor of the right thigh was taken from the patient with the help of the "loop" magnetic therapy device and the bioresonance therapy apparatus was recorded in 1 container. Immediately after that, globules from 1 container were transferred into container 3 (inversion), hand and foot electrodes were connected, the "loop" remained on the tumor area, the BRT session lasted 30 minutes. During the entire BRT session, the patient noted the characteristic sensations: tingling, "creeping", warmth in the area of the tumor of the thigh, chest. During all this time, the patient recalled the most stressful situations in his life as vividly as possible. After 30 minutes, globules were removed from 3 containers, destroyed in fire, "Information has been removed" from the containers of the apparatus. Then a clean crumble is placed in 1 container for recording. The recording was carried out for 10 minutes at the same positions of the "loop" and electrodes. During these 10 minutes, the patient tried to present his ideal, healthy image as vividly as possible. After the BRT session, the drug was tested through Cu met. D400, prescribed 6 globules 3 times a day for a month. After the BRT session, he underwent a course of chemotherapy according to the CAM scheme. A month later, 2 BRT sessions were performed similarly to 1 session, followed by chemotherapy according to the CAM scheme. The patient noted a continuous positive effect in the form of a decrease in the tumor of the hip and an improvement in clinical symptoms. After 2 months, the thigh tumor disappeared, there is no clinic, X-ray from The recording was carried out for 10 minutes at the same positions of the "loop" and electrodes. During these 10 minutes, the patient tried to present his ideal, healthy image as vividly as possible. After the BRT session, the drug was tested through Cu met. D400, prescribed 6 globules 3 times a day for a month. After the BRT session, he underwent a course of chemotherapy according to the CAM scheme. A month later, 2 BRT sessions were performed similarly to 1 session, followed by chemotherapy according to the CAM scheme. The patient noted a continuous positive effect in the form of a decrease in the tumor of the hip and an improvement in clinical symptoms. After 2 months, the thigh tumor disappeared, there is no clinic, X-ray from The recording was carried out for 10 minutes at the same positions of the "loop" and electrodes. During these 10 minutes, the patient tried to present his ideal, healthy image as vividly as possible. After the BRT session, the drug was tested through Cu met. D400, prescribed 6 globules 3 times a day for a month. After the BRT session, he underwent a course of chemotherapy according to the CAM scheme. A month later, 2 BRT sessions were performed similarly to 1 session, followed by chemotherapy according to the CAM scheme. The patient noted a continuous positive effect in the form of a decrease in the tumor of the hip and an improvement in clinical symptoms. After 2 months, the thigh tumor disappeared, there is no clinic, X-ray from prescribed 6 globules 3 times a day for a month. After the BRT session, he underwent a course of chemotherapy according to the CAM scheme. A month later, 2 BRT sessions were performed similarly to 1 session, followed by chemotherapy according to the CAM scheme. The patient noted a continuous positive effect in the form of a decrease in the tumor of the hip and an improvement in clinical symptoms. After 2 months, the thigh tumor disappeared, there is no clinic, X-ray from prescribed 6 globules 3 times a day for a month. After the BRT session, he underwent a course of chemotherapy according to the CAM scheme. A month later, 2 BRT sessions were performed similarly to 1 session, followed by chemotherapy according to the CAM scheme. The patient noted a continuous positive effect in the form of a decrease in the tumor o 10/07/98: there are no pathological changes in the lungs. The patient is observed in dynamics, as of March 2005, the state of complete remission.

2. Kh.D.A., born in 1973, map No. 2662. Registered since 09.01.01 with a diagnosis: stage III liposarcoma of the anterior abdominal wall. Histology: round cell liposarcoma with foci of myxomatosis.

Anterior abdominal wall represented by myself solid a crumbling purulent wound with a visible peritoneum in places. Due to the low effectiveness of chemotherapy in such a pathology, with the consent of the patient, 2 BRT sessions were performed similarly to the sessions in the patient from the previous example. During BRT, the patient noted similar sensory sensations. Immediately after the BRT sessions, chemotherapy was carried out according to the AR scheme. In total, 2 BRT sessions and 2 courses of chemotherapy were performed. After 2 months, a complete clinical cure was established; there was a thin scar on the skin of the anterior abdominal wall. As of March 2005, she is in complete remission. 3. P.E.P., born in 1939, map number 333. Registered since 03/27/98 with a diagnosis: stage II peripheral cancer of the upper lobe of the left lung. Sputum cytology: adenocarcinoma. Due to polyallergy and a mass of concomitant pathology, the operation is contraindicated. For palliative purposes, systemic chemotherapy was carried out according to the SR and CM regimens. A total of 5 courses were held. Due to the low expected effectiveness of chemotherapy, the patient received BRT sessions similar to the first two patients, situational homeopathy, which provided a stabilizing effect. For the last two years, the patient has been receiving constitutional drug Arsenicum, at first the 15th potency once every 2 weeks for 6 globules, then - the 25th potency once a month and now - the 25th potency once every 2 months. Chest X-ray: cicatricial changes in the upper lobe of the left lung, at the site of the projection of the tumor.

4. Sh.O.M., born in 1936, map No. 1369. Registered since 05.06.03 with a diagnosis of cancer larynx stage II. Histology: keratinizing squamous cell carcinoma. Due to age and concomitant cardiac pathology, the patient received palliative monochemotherapy with cyclophosphamide plus concentrations according to G.P. Grabovoi.

Task 1: the patient concentrated for 1-2 minutes on each digit of the laryngeal cancer code, mentally enclosing each digit in a transparent sphere and simultaneously imagining his ideal, healthy image inseparably with this digit. The number of concentration sessions is not limited. Since the patient, a retired military man, required an exact number of sessions, he was targeting 21 sessions per day.

Task 2: the patient, daily for 2 minutes, concentrated on the right index finger, mentally enclosing it in a transparent sphere and at the same time imagining his ideal, healthy image (from 2202 h to 2204 h).

3 task: daily from 2205 h 2210 h concentration on the selected color. Imagine pink, yellow, green, red, blue, purple colors in your mind. The color that appears brighter, stand out, represent within 5 minutes.

A month later, during an ENT examination, a tumor of the larynx was not found, in November 2005, the state of complete remission.

Conclusions:

1. Decisive for obtaining clinical recovery is

introduction in one way or another into the subconscious of the image of the patient's ideal health.

2. The patient's faith in the doctor is of great importance.

3. Information medicine methods are needed in the treatment cancer patients.

Gorbenko S.V. Clinical examples of curing cancer patients using information medicine methods // XI " IMEDIS ", 2005, vol. 2 -S.203-206