

Some aspects of the application of induction programs of brain rhythms
Kolganov I.V.
(Voronezh, Russia)

To break the closed cycle of stress - illness - stress, it is more effective to use induction programs (IP) not directly, but against the background of the load by the identified (s) indicator of the presence of a particular pathology.

We know that any diagnostic indicator (nosode, homeopathy, organopreparation, BR-preparation) is a medicine according to the principle of similarity. In the mode of electropuncture therapy, PIs can be the same medication.

Example

Patient M., 52 years old. From the anamnesis: considers himself ill for 2.5 years, which coincides with a divorce from his wife. Tested: Epiphysis D4, Adrenal gland D32, essential hypertension D30, herpes zoster virus D30-200, herpes simplex virus D12-60, candida glabrata, Intox I, duodenal ulcer D32, photon index - 17, etc. The optimal therapy step is determined through the herpes virus.

The sum of the potencies of the herpes virus ↓ + PI Development of self-esteem ↑, moreover, the intensity is 60 conv. units

The herpes virus is tested in the frequencies of the Stomach and Liver meridians. Hence FM drainage complexes were prescribed. During the session, it was proposed to retrospectively revive in memory the events of the last 3 years.

Reappointment in 1.5 months. Considers himself almost healthy. Intox II, herpes zoster, photon index - 13, pineal gland D4 are being tested.

Intox II ↓ + Epiphysis D4 ↑.

It can be assumed that the initial state at the time of the first visit is a recurring consequence. These pointers are tested in the frequencies of the meridians of the bladder, pancreas, and large intestine and are eliminated by PI Stress II.

At the third visit, Intox III, Epiphysis D4, D60 mushrooms (Candida glabrata) are tested in the frequencies of the gallbladder meridian. To conduct IP, the program Overcoming your fear is tested. The optimal photon index corresponds to the ideal one - 9. After the end of each IT session, the BR-preparation was recorded in the frequencies of all meridians in a simultaneous mode for 1-2 minutes, which was prescribed to the patient.

During the session, the patient experienced a psychological stress of 13-14 years associated with a quarrel between his parents. Three days later, his wife returned to him. Out of surprise, he called and asked for advice, to which he was offered to undergo another session of brain rhythm therapy.

Conclusion: selection of induction programs against the background of identified pointers or medications and a certain intensity makes the therapy more effective.

