

Statistical data obtained when working by the method of vegetative resonance test  
 "IMEDIS-TEST +" (results of work from May 2003 to February 2006)  
 G.Yu. Andreichenko  
 (Vorkuta, Russia)

In total, during this period, a four-level diagnosis of 686 people was carried out.  
 By age, the groups were distributed as follows.

Table 1

Age	Children under 1 of the year	1-10 years	11-20 years old	21-40 years old	41-60 years old	Over 60 years old	Total
Fact. count in	eight	105	99	189	252	33	686
%	1.1%	15.4%	14.4%	27.6%	36.7%	4.8%	100%

The current picture of the dominant miasms is presented in Table 2.

table 2

Dominant miasm	Fact. number of people	%
Botulism	6	0.9
Toxoplasma	five	0.7
Borrelia	3	0,4
Tuberculosis	33	4.8
Rickettsia	one	0.1
anthrax	one	0.1
Fungoid fungus	36	5.2
Coxsackie virus	five	0.7
Epstein-Barr virus	4	0.6
Cytomegalovirus	90	13.2
Hepatitis	43	6.3
Herpes	7	1.0
Rabies	3	0,4
Measles	7	1.0
Smallpox natural	10	1.5
Chicken pox	35	5.1
Polio	240	35.1
Rubella	138	20.2
Mycoplasma	2	0.3
Diphtheria	2	0.3
Syphilis	one	0.1
Pneumocystus	one	0.1
Psychovegetary load	13	1.9
TOTAL	686	100.0

The first 7 places in descending order are occupied by the dominant miasms of poliomyelitis viruses (35.1%), rubella (20.2%), cytomegalovirus (13.2%), hepatitis (6.3%), fungoid fungus (5.2%), chickenpox (5.1%), tuberculosis (4.8%). Which is 89.9% of the total.

In 13 cases, at the fourth level, only psycho-vegetative loads were tested, which were eliminated by a drug from the Bach Flowers group.

In 88 people, at the fourth level, both miasm and psycho-vegetative loads were determined, which were eliminated by Vipera 2000 in combination with a drug from the Bach Flowers group. Moreover, the Bach flower served in this case as a drainage.

Differentiating through the Intox II and Intox III pointers, the following picture emerged.

Table 3

Miasm	Intox II	Intox III	Total
Botulism	-	6	6
Toxoplasma	-	five	five
Borrelia	one	2	3
Tuberculosis	-	33	33
Rickettsia	-	one	one
anthrax	-	one	one
Fungoid fungus	-	36	36
Coxsackie virus	-	five	five
Epstein-Barr virus	-	4	4
Cytomegalovirus	3	87	90
Hepatitis	4	39	43
Herpes	one	6	7
Rabies	-	3	3
Measles	one	6	7
Smallpox natural	-	10	10
Chicken pox	five	thirty	35
Polio	five	235	240
Rubella	6	132	138
Mycoplasma	-	2	2
Diphtheria	-	2	2
Syphilis	-	one	one
Pneumocystus	-	one	one
TOTAL	26-3.9%	647-96.1%	673-100%

Analyzing the diagnostic picture, one can note a clear predominance of hereditary burden (96.1%).

For level 4 therapy, the following drugs were used: special.

Viper 2000 - 658 cases (95.9%); FM chords - 14 cases (2.1%);

nosode - 1 case (0.1%); Bach

flowers - 13 cases (1.9%).

After removing the miasms, I conduct therapy for deep mycoses.

Table 4 summarizes some data on the results of antifungal therapy. In total, at this stage, I observe 388 people. Directly there were 503 sessions.

The table does not include sessions in which work with allergens was carried out.

Table 4

Most commonly tested fungi

Type of fungus	First session	Repeated session	Total	%
Fungoid fungus	135	68	203	40.3
Dermatomycosis	59	6	65	12.9
Aspergillosis	35	4	39	7.8
Hansenella	13	-	13	2.6
Rodoturul	10	2	12	2.4
Pihia pini	13	2	fifteen	3.0
Saccharomycetes	eleven	one	12	2.4
Mukor pusilus	eight	-	eight	1.6
Schwaniomycetes	4	-	4	0.8
Actinomycosis	4	-	4	0.8
Candidiasis	2	-	2	0,4
Others (meetings. 1-3 times)	94	32	126	25.0
TOTAL	388	115	503	100 %

In subsequent stages antifungal therapy, the following began to emerge patterns.

Firstly, the regenerating complex RezoPLEX 19 began to be tested frequently.

the impression that it works at the stem cell level. Moreover, the body itself directs the action of the drug to the damaged tissue. The drug works for up to 2 months. It should be applied rationally after the functional systems of the cell have been restored, i.e. after miasm therapy, and when the process of renewal of protein structures is completed. After the preparation has been worked off, there is a further decrease in BI and PI, new reserves are opened up for bringing the body into a harmonious state.

Secondly, the pointer to allergic inflammations often began to work. Allergens in tissues, in this case, I regard as waste products of fungal colonies, accumulated during their long-term parasitism in the body. If you delve into the problem of allergens in tissues, then the conclusions are very serious. Allergens are even worse than free radicals because they are protein-based. And our DNA has a program to destroy someone else's protein - antibodies are formed. And if there are a lot of allergens in some tissue, the body begins to perceive this organ as a stranger, i.e. rejects it. Allergens cause fire on themselves and determine the general allergic readiness of the body.

#### Allergy symptoms

General: fatigue, decreased performance, sweating, chills, overweight, sleep disturbance, dizziness.

Psyche: decreased concentration, decreased memory, decreased mental performance, aggressiveness, irritability, depression.

Leather: eczema, urticaria, neurodermatitis, psoriasis, other lesions.

Digestive tract: nausea, heartburn, a feeling of fullness in the stomach, bloating, diarrhea or constipation, inflammation of the intestinal lining, Crohn's disease, ulcerative colitis.

Respiratory system: chronic rhinitis, sinusitis, tonsillitis, cough, chronic bronchitis, bronchial asthma.

Circulatory system: arrhythmias, tachycardia, high or low blood pressure.

Genitourinary system: hyperesthesia of the bladder, urinary incontinence, burning sensation during urination, frequent cystitis, diseases of the reproductive system in men and women.

Head: headache, migraine, conjunctivitis, otitis media.

Joints, muscles: swelling of the joints, pain in muscles and joints, rheumatic diseases.

Thirdly, after the complete recovery of the physical body, drugs for the correction of the psychological state began to be tested. In the selector "IMEDIS" there are very interesting drugs. Psychosocial loads "Medpharma". I use them very often in this case. I also test destructive psychological programs through a pointer to allergic inflammations.

I work through the pointer to the key problem (Fe met. D60) ↓ + the pointer to allergic inflammations (from the minimum test set) ↑ + OP (the organ with the most allergens) ↓ + medication ↑. The regimen for taking the drug is being worked out.

Thus, a scheme for testing and conducting stages of therapy in such a contingent of patients is being built.

---

G.Yu. Andreichenko Statistical data obtained by the method of vegetative resonance test "IMEDIS-TEST +" (results of work from May 2003 to February 2006) // XII

2006, vol. 1 - p. 297-302