

Bioresonance homeopathy - a differentiated approach to therapy
tobacco addiction

O.I. Speranskaya

(Federal State Institution "State Scientific Center for Social and Forensic Psychiatry"
Roszdrav, Moscow, Russia)

Tobacco dependence (TK) is one of the most common diseases in the world. At the same time, the effectiveness of TK treatment, despite the abundance and variety of existing methods, remains extremely low and amounts to no more than 25–40%, with the subsequent return of smoking in 80% of patients.

The use of bioresonance therapy can significantly (2–2.5 and more times) increase the effectiveness of treatment and prevent the occurrence of relapses of smoking. This is due to the possibility of effective use at given method treatment clinical diagnostic, clinico-pathogenetic approach, bearing in mind the neuroreceptor theory of TK and pathogenesis knowledge of the clinical forms of the disease.

We used exogenous bioresonance therapy fixed frequencies on devices "MINI-EXPERT-DT" and endogenous bioresonance therapy on the device "IMEDIS-BRT-A" (circular or diagonal BRT) with additional connection of cassettes with homeopathic medicines.

So, with the ideator form of TK, the most preferable is the use of the dopaminergic program (E 34).

In the dissociated form of the disease, therapy should be started with activation of cholinergic receptors (program E 67), then you can recommend dopaminergic program followed by a session of BRT.

With the most severe and complex for treatment In the psychosomatic form of TK, the most effective is the sequential application of programs for the activation of adrenergic, cholinergic and dopaminergic systems (E 186, E 67, E 34) and BRT sessions.

Electro-acupuncture diagnostics, along with clinical studies, allows you to track how the patient's neurohormonal background, his mental state, craving for tobacco smoking, the state of autonomic regulation of functions during therapy and at the end of the treatment session are changing.

In the intervals between sessions, patients are recommended to use resonance homeopathy, individually selected testing of on results hormonal levels or electronic programs (E 34, E 67, E 186, zoned copies respectively).

The proposed method of treating patients with various clinical variants of TK syndrome allows you to refuse to take funds psychopharmacotherapy, which allows it to be widely used in outpatient practice, including for the treatment of children and adolescents.