Management of patients with chronic diseases of the joints and spine using BRT and homeopathy E.A. Matyushenko, O. I. Eliseeva (LLC "Eliseeva Methodological Center", Moscow, Russia)

The problem of chronic diseases of the joints and spine, such as rheumatoid arthritis, psoriatic arthropathy, ankylosing spondylitis, articular form of rheumatism, gout and other nosological forms, accompanied by joint damage, is becoming more and more relevant every year, especially among the female population. There are two concerns about it.- on the one hand, the number of young patients who fall ill during adolescence and adolescence increases, on the other hand, articular syndrome begins during menopause against the background of age-related osteoporosis and a decrease in the level of estrogen in the blood.

Against the background of the increasing threat of disability of patients and a large number of side effects caused by traditional therapy sulfonamides, hormonal drugs and cytostatics, there is a need to search for new alternative ways of managing such patients. Under these conditions, the effectiveness of bioresonance therapy (BRT) methods in the management of rheumatic patients is difficult to overestimate.

For the analysis, the results of the examination of 22 patients over 2 years were used, of which 18 - women between the ages of 25 and 70. All of them had pain in the area of large and small joints, swelling of the joints and limited mobility, two of them had flexion contractures.

As you know, the basis for the onset of chronic inflammation in the synovial tissue of the joints is the state of long-term immunodeficiency resulting from the persistence of a chronic viral infection. Most often in this group of patients, a combination of herpes viruses of 1 and (or) 2 types, Epstein-Barr virus and beta-hemolytic streptococcus was tested. Most often, one of the two above-mentioned viruses is tested in the D15 potency, the other in the D30 potency, streptococcus is detected in the D8 potency

up to D15 depending on the severity of the process. Somewhat less meets often, chlamydial joint damage, most often in the D15 potency. All x patients tested nosodes of infectious arthritis, rheumatism and nosodes testing and treatment of rheumatoid diseases. Depending on the severity of the process, their potencies ranged from D6 to D15. In addition, all patients were tested with suitable organ remedies for the affected joints, most often in the D6 potency. This fact usually implies the active use of frequency therapy, nosodes and organopreparations in treatment, however, practice has shown that their active use often provokes an exacerbation of the articular syndrome, which sharply reduces both the patient's quality of life and his faith in treatment. This required the search for more environmentally friendly therapy methods for the patient and the use of complex techniques. We have worked out the following recommendations:

1. Frequency therapy is undesirable to carry out at once on all identified microorganisms, because this provokes an exacerbation. It is wiser to conduct sessions only at one specific frequency, it is better to start with the bacterial frequency, which is identified as the cause of the inflammation. At the same time it is advisable

to carry out a standard course of antiparasitic therapy and a course of cleansing the body, which includes tubing of the liver and irrigation of the intestines according to the patented method of O.I. Eliseeva. Cleansing the body is very effective for metabolic diseases of the joints and lungs,

compensated course of autoimmune polyarthritis. However, the appointment of a classic cleansing course should be treated with caution, because fasting and tubing can cause exacerbation, and therefore are contraindicated in patients with decompensated arthritis. In such cases, it makes sense to test and prescribe Hepar Comp to ensure better drainage of the liver. ("Heel") 10 intramuscular injections-1 time in 4-5 days. In addition, a natural immunomodulator must be prescribed for patients with a chronic decompensated course of arthritis.- Transfer factor (TF). For viral etiology, we prescribe the Transfer Factor Advensd, 6 capsules per day, with the autoimmune process TF, the classic is prescribed for 4-6 capsules a day. It was noted that against the background of the use of TF, the cure terms of the process are reduced, and the exacerbation, if it occurs, is in weak manifestations. In combination with TF, the treatment result is achieved more effectively.

2. Technique of BRT. The therapy is carried out according to the tested meridians (most often with polyarthritis these are the meridians of the spleen and pancreas, articular degeneration, connective tissue

degeneration and liver), the inductor is connected to the frontal electrodes and placed on the most affected joint, clean crumbs are placed in the first container, and BRT is performed for 1 second for each meridian (the total therapy time should not exceed 10 seconds). After that, the crumbs from the first container are rearranged into the third, and the tested nosodes and organopreparations are connected from the selector. It is important to note that all drugs chosen for therapy must be either in the same potency (with potency D30 and higher), or they can be in different potencies (if the potency is from D5 to D15). For the best effect, it is necessary to use drainage, and in this case it makes sense to test and connect at the same time resoplexes from the Schimmel group of drugs (lymph, pancreas, stomach, liver, kidneys, female endocrine system, paranasal sinuses, inflammation). BRT is carried out for at least 20 minutes, after which the selector is turned off, and in the first container, the preparation is written on clean crumbs for 3 minutes, while the crumbs in the third container remain in place. Now the drug is ready, its dosage is individually tested. It turns out that we are not recording the drug itself, but the body's reaction to it, which arises against the background of pathological oscillations set in inversion. This method of recording provides a milder therapeutic effect with less likelihood of exacerbation, and if it does occur, it is recommended to take a break 3 that we are not recording the drug itself, but the body's response to it, which arises against the background of pathological oscillations set in inversion. This method of recording provides a milder therapeutic effect with less likelihood of exacerbation, and if it does occur, it is recommended to take a break 3 that we are not recording the drug itself, but the body's response to it, which arises against the background of pathological oscillations set in inversion. This method of recording provides a milder therapeutic effect with less likelihood of exacerbation, and if it does occur, it is recommended to take a break 3-4 days before the symptoms subside, then switch to a dosage of 1 grain 1 time per day. This drug is conventionally called BR # 1.

Another preparation is prepared using a time modulation mode. To do this, either the previously selected meridians are left, or all meridians are turned on, after which we find the "effects" section in the time modulation and the frequencies are tested. The most important here are anti-inflammatory, analgesic, arthrogenic, vertebrogenic, anti-gout, antioxidant, geopathogenic, immunostimulating effects, the frequency of the bacterial pathogen can also be added to them, which is an inducer of the inflammatory process (best of all - one frequency). Clean crumbs are placed in the first container and BRT is carried out. Treatment time 7-10 minutes. The drug is code-named BR # 2. The rules for dose selection and administration are the same.

3. Against the background of ongoing therapy, it is recommended to include homeopathy in in the form of chelators - a complex of Heel preparations: Traumeel C, Zeel T, Diskus Comp in 10 intramuscular injections-15 injections, alternating with an interval of 1 to 3-4 days between injections, depending on the severity of the process, - the sharper the current, the more often. Their use causes a pronounced positive trend,- swelling and stiffness of the joints and spine go away, pains significantly decrease and even disappear, patients note a significant influx of strength, both physical and moral. If Hepar Comp. Was used, then the injections of these drugs are started after the end of the Hepar Comp. Course. It is also recommended to use Traumeel C ointment for local action. In cases of prolonged pain syndrome, we carried out injections around the joint with Traumeel C preparation daily 5-7 injections per course. Such a comprehensive treatment leads to the removal of swelling, inflammation and pain in the joints.

4. It should be remembered that during treatment we must support the level of quality of life of patients - for this it is advisable to use NSAIDs, and it is better not in tablet form (since they cause damage to the stomach), but in suppositories.

5. In 8 out of 10 patients, the polarity was reversed. Most an effective way to eliminate it is the method outlined by O.I. Eliseeva at the XII International Conference.

1) A bioresonance preparation BR-1 is being prepared according to the method of A.A. Hovsepyan.

2) Testing and correction in the CNS section of the frequencies of rhythm disturbances brains, program E, intensity L-15, magnetic "belt" placed on the spine, "loop" on the cervicothoracic spine, -

duration - 3-7 minutes per frequency, 3 in total-5 sessions, every day or every other day. In addition, Systemic Spiritual Adapters are used. Recommended scheme for the use of Systemic Spiritual Adapters- selection 1-2 from the recommended list, after which they are assigned 1 grain of one specific type 1 time per day before bedtime. Many patients note, against the background of their reception, the appearance of specific dreams that affect their spiritual state. It should be noted that the pathological informational basis of these diseases arises not only because of the social negativity acquired by the patient himself in the process of life, but it is also associated with the information that the patient receives from his kind, and here it makes sense to talk not only about molecular genetic aspect of the etiology of these serious chronic diseases, but also about deeper spiritual moments associated with the moral state of both the patient himself and his ancestors. The presence of social negative in the ancestral past of the patient (which, unfortunately, is very possible,

6. When carrying out diagnostics, it is important to pay attention to the presence radiation to organs such as the thyroid gland, adrenal glands and bones.

Radiation is removed by preparations of Radium, Strontium, Plutonium, Uranium, by Imedis Anti-Radiation Comp., By preparations by Rayex.

7. Strengthening of the articular syndrome occurs against the background of a decrease in production estrogens, either against the background of menopause (6 patients), or ovarian dysfunction (2 patients). Correction of these conditions should be carried out by the joint efforts of a gynecologist-endocrinologist with the help of hormonal drugs and a homeopath. Here it will be appropriate to use the preparations Hormel and Klimakt from Heel. In 3 patients, the autoimmune process in the joints was combined with autoimmune thyroiditis. Here, in order to achieve stable compensation, it is necessary to correct the hormonal background, which is achieved both by the use of hormonal preparations, and again by the use of homeopathy.

- preparation Strumel from Heel.

8. When testing, pay attention to the presence of geopathogenic and electromagnetic loads in the body. The geopathogenic load was tested in 16 of 22 patients. The elimination of the geopathogenic load was carried out according to the scheme adopted in the methodological center of O.I. Eliseeva:

1) Homeopathic crumbs with a frequency of 6.2 Hz, which is prescribed in 3 mountains. 2 times a day for 2-4 weeks.

2) It is recommended to rearrange the bed in the apartment. Place in the apartment determined by dowsing.

3) Carry a Gamma-7 device with you, which protects against negative loads - geopathogenic and electromagnetic.

Conclusions:

1. When starting the treatment of autoimmune joint disease, it is necessary explain to the patient that this treatment will be long and will require a lot of patience and work, not only in fulfilling all the recommendations, but above all in working on oneself and one's perception of life. Only in this case it is possible to achieve a stable, until the end of life, remission of the disease.

2. A comprehensive method of treatment using BRT, homeopathy, methods A.A. Hovsepyan, O. I. Eliseeva, against the background of the use of Transfer Factors, makes it possible to achieve 100% tangible shifts in treatment and longterm remission in chronic diseases of the joints and spine.

E.A. Matyushenko, O. I. Eliseeva Management of patients with chronic diseases of the joints and spine using BRT and homeopathy // XIV "- M .:" IMEDIS ", 2008, v.1 - C.206-212