

Statistical data on the use of multilevel systemic therapy in the joint medical practice of MCIT "Artemis" and GUZ "TsVMiR No. 1" RO for 2007

A.E. Kudaev, N.K. Khodareva, S.V. Kruglova

(Medical Center for Innovative Technologies "Artemida", Rostov-on-Don, Russia)

The results of the application of the method of multilevel systemic therapy [1] in the clinical practice of MCIT "Artemis" and GUZ "TsVMiR No. 1" RO, for 2007 are shown in Table 1.

Table 1

Nosology	Rack. improved (number sick.)	Relates. improved (number sick.)	Without. improved (number sick.)	Total (number sick.)
Ill. respiratory organs	58	fourteen	-	72
Diseases of the cardiovascular. systems	49	61	-	110
System diseases digestion	215	85	2	302
Diseases on-check and urinary system Diseases of the central	142	38	-	180
nervous system and peripheral nervous systems	12	sixteen	4	32
Diseases of the support-but- locomotor apparatus	24	18	-	42
Diseases of the skin and hair	29	39	eight	76
Diseases of the genital area	243	67	-	310
Diseases of the endocrine system	18	43	4	65
Fate science	75	70	2	147
Total	865	451	twenty	1336
Total, %	64.76	33.8	1.5	

All patients listed in this table were taken for treatment after unsuccessful therapy by methods of classical medicine.

Under lasting improvement either complete recovery or the onset of stable remission during a chronic disease, confirmed by the data of an objective clinical examination, was understood. Under relative improvement the improvement of the general condition, the transition of the disease to a subacute state, accompanied by the presence of the main symptoms, was understood.

One of the most pressing problems nowadays is female infertility. In our center there is a proprietary methodology that allows using the equipment of the Center "IMEDIS" to effectively and in a relatively short time to solve this problem. We bring to your attention one of the clinical cases.

B-I Anna M., 35 years old. Turned for help in the MCIT "Artemis" 15.05. 2007
Complaints about the inability to get pregnant during the last three years; tense relationship with her husband, inability to find a compromise in solving vital problems. Has been married for 15 years; a history of 3 honey. abortion, last in 2002, without complications. Regular sex life without contraception. At the time of examination, the psychoemotional state was unstable; pronounced asthenic syndrome; worried about insomnia, decreased performance.

In the period from 2005 to 2007, she was repeatedly treated in a hospital and on an outpatient basis for the inflammatory process of the uterine appendages. According to the results of PCR diagnostics, chlamydia, Trichomonas, ureaplasma, gardnerella were identified. Several courses of antibiotic therapy were carried out to both partners - without effect. Spermogram of the husband without pathological changes; postcoital test - 100% sperm death in the vagina.

Testing revealed:

- total radioactive load of the 2nd degree;
- geopathogenic load of the 2nd degree;
- low reserves of adaptation of the 4th degree;
- stress of the endocrine system of the 2nd degree;
- hormonal imbalance;
- signs of chronic salpingitis of bacterial etiology. Both partners underwent examination and treatment at MCIT "Artemis".

After carrying out the technique of adaptation of the patient to her husband for three weeks, family relationships improved significantly; decreased conflict; there was a mutual desire to discuss and solve current problems. Along with this, the patient clearly developed a desire to have a child from her husband, and not "to have a child in general".

Against the background of treatment with targeted nosodes, the revealed genital infections were completely eliminated within two months; PCR diagnostics - negative result.

In the course of therapy, the following nosodes were used:

- inverse targeted blood autosode;
- inverse targeted autosode of discharge from the patient's genital tract;
- inverse nosode of the amount of genital infections;
- the inverse sperm nosode of the husband.

In the course of therapy, chronosemantics was performed twice - with standard CSP and with an inverse nosode of discharge from the genital tract adapted to the CMH. In addition, the cycle of therapy included the correction of psychosomatic disorders with drugs "Medpharma", "Rubimed", "Bach Flowers". In October 2007, she was adapted to her husband's sperm; in November 2007 - targeting the husband's inverse sperm nosode to vaginal discharge. 12/22/2007 is the date of the last normal menstruation. 19.02. In 2008, a normal uterine pregnancy was diagnosed for a period of 6-7 weeks. At this stage, the pregnancy proceeds without pathology. The total duration of therapy since the initial visit was 8 months.

Thus, systemic multilevel therapy is successfully used in the treatment of mixed infertility.

Literature

1. Kudaev A.E., Mkhitarian K.N., Khodareva N.K. Multilevel systemic therapy targeted energy-informational preparations and Systemic Spiritual Adapters. - T.: LLC "Publishing house Lukomorye", 2005. - 128 p.

A.E. Kudaev, N.K. Khodareva, S.V. Kruglova Statistical data on the use of multilevel systemic therapy in the joint medical practice of MCIT "Artemis" and GUZ "TsVMiR No. 1" RO for 2007 // XIV

"- M.:" IMEDIS ", 2008, vol. 2 - p. 24-27