

The method of preparing a targeted urine autonosode in combination with general bioresonance drug

V.A. Brichuk, O. V. Koritsky
(PE "Medicor-2", Vinnitsa, Ukraine)

Long-term practice of using urine autonosode and OBR- and PBS preparations prompted the idea of combining these techniques in various combinations.

For these purposes, a stationary computer complex "IMEDIS-EXPERT" manufactured by LLC "IMEDIS", a device for autonomous BR-therapy "IMEDIS-BRT-A" and an autonomous drug selector are used.

Preparation of the drug begins with the recording of the KMX marker. For this, the chronosemantic light probe (designed by Kudaev, Mkhitarian, Khodareva) is connected to the socket of the container No. 2 of the IMEDIS-BRT-A apparatus. Place a glass of homeopathic crumbs in container # 1 and record information from the main chiroglyphic lines of the hand (Life Line, Destiny Line, Head Line and Heart Line) for 60 seconds from each palm of the patient. Then, in container No. 1 of the "IMEDIS-BRT-A" apparatus, we record pathological oscillations in the automatic BRT mode along all meridians for 60 seconds. Then we move the glass with crumbs from cotneyer No. 1 into container No. 3 (inversion).

After that, in the drug selector, we select a number of the necessary organopreparations in the D6 potency and a number of drugs of the DRE and DIS groups of the ONOM company, depending on the patient's diagnosis. In the 2nd container of the "IMEDIS-BRT-A" apparatus is placed a previously prepared and transcribed into crumbs autonosode of urine of the given patient in D12. We carry out the potentiation of the autonosode when the Medication Selector apparatus is connected to the container No. 2. Place the KMX marker in the passive electrode in the patient's hand.

We potentiate the autonosode by rotating the transfer knob from 7 to 0. When potentiating, the following condition must be met:

KMX marker + Selected potency of urine autonosode .

We put a cup with crumbs in the 1st container and record the combined preparation for 15 minutes, in the automatic BRT mode, a "belt" device is connected to the frontal electrodes, which is thrown over the patient.

At the end of this time, the preparation is ready. Inversion and load is removed. An individual dose is selected according to the KMX marker:

KMX marker + Increasing number of grains of the combination preparation

.

The resulting drug is prescribed once a day, in the evening.

For a deeper and longer action of the drug, a conventional chronosemantic probe is connected to the socket of container No. 2, which is applied to the end point on the patient's Life Line (for men - on the left hand, for women - on the right) and in this position the combined drug is recorded during 15 minutes.

This technique allows you to deeply and long-term on various pathological processes, to manage them. You can use nosodes of various environments of the human body and adjust their action.

This technique has been used for the last 2 years in 38 emergency patients "Medicor-2" (Vinnytsia) with the following nosologies:

- widespread osteochondrosis of the spine, multiple vertebral hernia - 23,
- gout, gouty mono and polyarthritis - 7,
- rheumatoid arthritis - 3,
- viral hepatitis B and C - 5.

The course of treatment took from 6 months to 1.5 years, all patients managed to achieve a stable remission of the disease, everyone continues treatment with pleasure. This technique can be recommended for the complex, long-term treatment of metabolic diseases and autoimmune diseases.

Literature

1. Gotovsky Yu.V., Bobrov I.A., Goltsov A.A., Kosareva L.B., Mkhitarian K.N. Virtual space of internal nosology of the body and its compactness // Abstracts and reports of the XI International conference "Theoretical and clinical aspects of bioresonance and multiresonance therapy". - M.: IMEDIS, 2005.
2. Kudaev A.E., Mkhitarian K.N., Khodareva N.K. Systemic spiritual adapters and their role in modern energy-informational medicine. // Abstracts and reports of the XI International conference "Theoretical and clinical aspects of bioresonance and multiresonance therapy". - M.: IMEDIS, 2005.
3. Kudaev A.E., Mkhitarian K.N., Khodareva N.K. Multilevel systems diagnostics of the body using resonance chains within the framework of virtual ART // Ibid. - M.: IMEDIS, 2005.
4. Gotovsky Yu.V., Mkhitarian K.N. Chronosemantic diagnostics and mantic point therapy. 2nd ed. rev. and add. - M.: IMEDIS, 2002. --- 392 p.
5. Gotovsky Yu.V., Mkhitarian K.N. Lectures on chronosemantics. - M.: IMEDIS, 2004. --- 276 p.
6. Kudaev A.E., Mkhitarian K.N., Khodareva N.K. Multilevel system therapy with targeted energy-information drugs and Systemic Spiritual Adapters. - T.: LLC "Publishing house" Lukomorje, 2005. - 128 p.
6. Koritskiy O.V., Koritskiy Yu.V. Application of targeted autonomodes urine and blood for the treatment of chronic urethrostatis // Abstracts and reports of the XII International conference "Theoretical and clinical aspects of bioresonance and multiresonance therapy". - M.: IMEDIS, 2006.

V.A. Brichuk, O. V. Koritskiy Method of preparing a targeted urine autonomode in combination with a general bioresonance drug // "M.: IMEDIS", 2008, vol. 2 - P.79-82