

The use of endogenous bioresonance therapy with moving magnetic electrodes

O.A. Postnikova, E.V. Postnikova

(Department of Homeopathy and Electropuncture IPK FMBA, Moscow, Russia)

The authors propose a method for using bioresonance therapy with additional magnetic inductors that move from the head to the lower body procedure time in a certain sequence.

The procedure is carried out against the background of a general BRT with cylindrical and (or) with flat electrodes, or with point electrodes located on biologically active points along the key meridians with testing the optimal location of the electrodes on the patient's extremities. The placement of the main electrodes of general action on biologically active points accelerates the reaction time to the effect of BRT, reduces the time of the procedure and makes the effect as accurate and directed as possible to the solution of the problem. Therefore, we give preference to BRT with a choice of acupuncture points in accordance with the nature of the pathology and the patient's personality.

Additional electrodes are located on the zones having maximum deviation from the norm, increase the weight, the importance of these zones for assessment by the body, which allows the body to focus its efforts on the problem emphasized by the procedure. The use of magnetic inductors as additional therapeutic electrodes allows you to remove information and carry out the effect not on the surface of the skin, but at a deeper level due to their magnetic effect. It is important to position the electrodes so that the diseased organ or tissue is between them. It is possible to connect one inductor to the affected area and then move it to the contralateral side or from the front to the back with alternating both positions, or just the places to be treated.

In many cases, to restore general vegetative balance, the inductor must first be located in the face, most often in the mouth, then move to the back of the head and then its position should alternate: either on the front surface of the body, then on the back. In this case, the level of exposure falls down gradually, reaching the tailbone and the lower end of the body. For problems in the upper and lower extremities, they should also be involved in therapy.

The mouth area is especially important, as it is the most prominent part of the body in front. If we consider the embryonic period (and a person in phylogenesis), then we can conclude that the most ancient or earliest way of exploring the world is the method of contact of the object under study with the tongue - taking it into the oral cavity. In the course of therapy, the optimal position of the electrode is tested at each current moment, and the body dictates the mouth as the first zone for activating life processes with the help of BRT, revitalizing the processes of interaction between the body and the environment.

In case of gross pathological processes, the introduction of pathological information from the body into the resonant process in an unchanged form or reinforced by an inductor is undesirable. In this case, the signal from the source must first be

weakened or inverted. And then, too, depending on the depth of the pathological focus under the skin, a simple electrode or magnetic inductor should be used. However, you should monitor the duration of exposure to the inverse electrode and in time switch to direct exposure, when the disharmonic process weakens, it is better to first use the zones around the focus.

In the work, with the correct choice of the sequence and method of action, there is an improvement in well-being during the procedure, pain passes, "enlightenment in the head" appears, and so on. During the treatment, there is an improvement in the ART of the psycho-emotional state, the level of health, there is an optimization of resources and other indicators of ART (according to the scales developed by the authors). In case of suboptimal selection of the treatment zone, discomfort may appear, which can be corrected by changing the placement of the electrodes.

It is important to change the strength of the BRT signal using the potentiometer knob, adjustable depending on the tolerance of therapy and the capabilities of the body: from strong to very weak, which is actually capable of a deeper effect and gives a long-term effect. However, such weak signals should not be applied immediately at the beginning of treatment, this effect is similar to the use of high potencies in homeopathy, and the body should be prepared for it.

An example is the treatment of BRT in this modification: there was a resorption of hemorrhage into the vitreous body, into the retina, the shape of the deformed eyeball was aligned with astigmatism, myopia, and the mobility of the eyeballs was restored with strabismus. In these cases, one electrode was located in the area of the eyeball, the other on the occipital area. The signal intensity also changed during the procedure.

The use of moving electrodes on the body allows you to quickly relieve pain in the spine, in the gallbladder, spastic phenomena in the intestines, symptoms of vegetative dystonia, etc. and at the same time avoid suppressing symptoms, help normalize the activity of the chakras, relieve congestion in certain areas of the body and promote the integration of individual parts of the body into a single whole (including the head and body), mental and bodily dynamic processes.

O.A. Postnikova, E.V. Postnikova Application of endogenous bioresonance therapy with moving magnetic electrodes //

"IMEDIS", 2008, v.2 - C.289-292