

## Possibilities of the modified method of electropunctural diagnostics in determining the mutual compatibility of homeopathic medicines

G.A. Yusupov  
(JSC Holding "EDAS", Moscow)

One of the unresolved issues in homeopathy is the possibility of prescribing several homeopathic remedies together. It is known that even the founder of homeopathy, S. Hahnemann, often in cases of joint use of two or more homeopathic remedies noted the absence or decrease in the effectiveness of treatment, and with the individual use of each of them, the therapeutic effect was expressed to one degree or another. At the same time, there is a lot of evidence of convincing results when using complex homeopathic preparations produced by Heel, EDAS, etc. The practical experience of many generations of homeopaths also indicates that some homeopathic preparations, when used together, have a full therapeutic effect, often complementing each other, and the combination of some drugs mutually reduce the effectiveness of the treatment. In the work of R. Gibson Miller provides data on the mutual incompatibility of more than 60 known homeopathic remedies [1]. Revealing mutual compatibility requires long-term practical experience, a scrupulous analysis of the results of treatment with the combined use of several homeopathic drugs and comparing these results with the effectiveness of therapy with one of the drugs. Such studies are laborious, time-consuming and, therefore, in the scientific literature, one can rarely find works devoted to this topic. a scrupulous analysis of the results of treatment with the combined use of several homeopathic medicines and comparing these results with the effectiveness of therapy with one of the drugs. Such studies are laborious, time-consuming and, therefore, in the scientific literature, one can rarely find works devoted to this topic. a scrupulous analysis of the results of treatment with the combined use of several homeopathic medicines and comparing these results with the effectiveness of therapy with one of the drugs. Such studies are laborious, time-consuming and, therefore, in the scientific literature, one can rarely find works devoted to this topic.

The aim of this work is to study the possibility of a modified method of electro-acupuncture (energy-informational) diagnostics with drug testing in determining the mutual compatibility of homeopathic preparations when used together.

If you fully follow the method of electropunctural diagnostics with drug testing according to R. Voll, it is necessary to start diagnostics with quadrant measurements: "hand - arm", "right arm - right leg", "left arm - left leg" and "leg - leg". This measurement uses two hand-held tubular electrodes and two brass plate foot electrodes. An indicator equal to 82–84 USD is considered to be normal. Indicators above the norm indicate an excess of the energy balance in the corresponding part of the body, while indicators below the norm indicate a decrease in the energy balance. In cases when the quadrant measurements are below 80, it is necessary to "recharge" - a weak current is passed through the hand electrodes with a smoothly varying frequency range from 0.9 to 10 Hz and vice versa, the so-called "swing".

After quadrant measurements, a study of electrical conductivity (EC) is carried out at the control points of all 20 meridians. On the meridian with the greatest deviations of the EP indices at the control point, the indices of other points are examined. Diagnostic conclusions are made on the basis of the indicators of the dynamics of the indicators of the measuring device. Great importance is attached to the speed of movement and the parameters of the "arrow fall". In the tactics of treatment, much attention is paid to "foci of burden" and the relationship of organs with the corresponding teeth. Homeopathic treatment is selected by drug testing. In total, the examination of one patient and treatment procedures according to the Voll method take from 1.5 to 2.5 hours [6, 7].

The advantages of the modified method [2] are:

1. Reducing the duration of the procedure to 15–20 minutes. Study of the dynamics of EP carried out in 5-7 biologically active points (BAP) located only on the fingers.
2. Simplicity and consistency of interpretation of diagnostic results. Dynamics is assessed

electrical conductivity before and during modulation of the measuring current by the wave characteristics of homeopathic preparations. There is no need to take into account the "arrow falling" phenomenon. At the beginning, the EP is measured without connecting the tested drugs, then each drug is tested separately and at the next stage joint testing is carried out. If, during separate and joint testing, EP indicators improved compared with the initial ones, the drugs are compatible. If during separate testing the EP improved, and during joint testing, on the contrary, the indicators worsened, the tested means are not compatible.

During the testing of homeopathic preparations, the mutual incompatibility of which is known from the literature data, a deterioration in the indicators of electrical conductivity was noted when they were jointly tested. With separate testing of each of them, the EP indices improved to varying degrees. Similar results were obtained during testing using the wave characteristics of homeopathic preparations from a drug selector, which was a ROM microcircuit. The hardware and software complex allows you to quickly connect the wave characteristics of drugs for testing in various combinations to the measuring current and evaluate the expected efficiency of their combined use by the dynamics of the EP. Using these capabilities of the method, during the outpatient appointment, it was possible to accumulate material, on the basis of which the mutual incompatibility of 44 more homeopathic preparations was determined (Table 1). A feature of the study is that the identification of incompatibility was possible only in cases when a patient is examined, in whom testing separately for each drug improves the EP BAP, and when they are simultaneously connected, on the contrary, it worsens. Disabling one of the drugs again leads to an improvement in the BAP electrical conductivity. The final decision on the mutual incompatibility of certain homeopathic remedies was made after a similar dynamics of electrical conductivity was observed in at least 10 patients. At the same time, there should not be a single case when joint testing did not lead to a deterioration in the EP BAP indicator [2, 4].

Table 1

N / a	Name drug	The serial number of the incompatible drug						
1	Acidum f luoricum	38						
2	Aconitum napellius	12	22	thirty	32	40		
3	Adonis vernalis	35						
4	Aesculus	23	thirty					
5	Aloe	22	35	37	40			
b	Anacardium	ten	25	33				
7	Apis mellifolium	eight	41	42				
eight	Arnica montana	7	22	23	28	33	36	
nine	Baptisia	16	24					
ten	Barium jodatium	b						
eleven	Belladonna	28	32					
12	Berberis	nine	24	35	37	38	41	
13	Brionia	19	26	32	37	33		
fourteen	Cactus	fourteen	thirty	36				
15	Calcium carbonicum	13	22	thirty	32	35		
16	Capsicum	nine	43					
17	Causticum	eighteen	23	35				
eighteen	Cinnabaris	17	twenty	22	34	39	42	
19	Collocynthis	13						

twenty	Drosera	eighteen	33	34				
21	Gelsemium	eight	35					
22	Graphites	2	5	15	eighteen	thirty	41	
23	Hamomellis	eight	17	27				
24	Hydrastis	nine	12					
25	Ignatia amara	eight	thirty	32				
26	Kalium bichromicum	28	32					
27	Lapis album	23						
28	Mercurius solubilis	eight	26	35	37	38	40	
thirty	Nux vomica	2	fourteen	15	22	25	32	41
31	Phosphor	28	thirty	32	38			
32	Pulsatilla	2	eleven	13	15	25	thirty	31
33	Rhus toxcodendron	2	b	eight	13	twenty	22	44
34	Sarsaparilla	eighteen	twenty	33				
35	Scrophullaria	3	5	12	15	17	21	28
36	Secale cornutum	eight	fourteen	39				
37	Sepia	5	12	13	28	31		
38	Silicea	1	31	12	28	40		
39	Solidago	eighteen	44					
40	Sulfur	2	5	28	38			
41	Thuja	12	22	thirty				
42	Urtica urens	7	12	eighteen	43			
43	Veratrum album	2	16					
44	Zincum metallicum	thirty	33	39				

Since the results of treatment with drugs or their energy-informational analogues, testing of which normalizes electrical conductivity is significantly higher than from the results when taking placebo, it can be concluded that the dynamics of EP BAP indicates the compatibility or incompatibility of various homeopathic drugs. If the drugs have not revealed mutual incompatibility, then the joint appointment of several homeopathic drugs is quite acceptable. For example, if a patient with peptic ulcer disease with pain syndrome has heartburn and signs of bleeding, it is difficult to find one homeopathic remedy that covers all manifestations. The combined use of compatible drugs: Anacardium, Collocynthis, Hamomellis and Hydrastis has a full therapeutic effect in 87% of patients with duodenal ulcer disease. Additional use of the Helicobacter pylori nosode leads to the eradication of bacteria in more than 90% of cases [5]. It is important to note that joint testing of these drugs or their electronic analogs leads to the normalization of EP BAP values also in about 90% of patients with an established diagnosis of peptic ulcer disease. In cases where heartburn is not noted, the dynamics of EP BAP is not observed when testing the Hydrastis drug, and if there is no bleeding from the ulcer, there is no BAP response to the energy-informational characteristics of the Hamomellis drug. Our studies have shown that the results of treatment with mono-drugs, taking them together, or a complex drug, which includes these components, as well as their electronic copies, are identical [2, 3, 4]. that joint testing of these drugs or their electronic analogs leads to the normalization of EP BAP values also in about 90% of patients with an established diagnosis of peptic ulcer disease. In cases where heartburn is not noted, the dynamics of EP BAP is not observed when testing the Hydrastis drug, and if there is no bleeding from the ulcer, there is no BAP response to the energy-informational characteristics of the Hamomellis drug. Our studies have shown that the results of treatment with mono-drugs, taking them together, or a complex drug, which includes these components, as well as their electronic copies, are identical [2, 3, 4]. when heartburn is not noted, the dynamics of EP BAP is not observed when testing the Hydrastis preparation, and if there is no bleeding from the ulcer, there is no BAP response to the energy-informational characteristics of the Hamomellis preparation. Our studies have shown that the results of treatment with mono-drugs, taking them together, or a complex drug, which includes these components, as well as their electronic copies, are identical [2, 3, 4].

Based on the results obtained, it can be assumed that the recommendations of the founder of homeopathy S. Hahnemann to prescribe only one drug were due to the mutual incompatibility of individual drugs. Clarification of specific cases of incompatibility by the method of electropunctural diagnostics with joint drug testing, confirmed by clinical studies, will avoid their combined use in medical practice.

The use in a complex of preparations of compatible and complementary or ready-made complex homeopathic medicines facilitates the selection of homeopathic treatment and increases efficiency. When developing new complex drugs, it is necessary to take into account the mutual compatibility of their components.

#### Literature

1. Gibson R. Miller. Relationship and duration of action of homeopathic drugs. - Smolensk: Homeopathic Medicine, 1994.
2. Yusupov G.A., Zilov VT., Khadartsev A.A. Modified technique electropuncture (energy-informational) diagnostics in practical medicine. A guide for doctors. - M., 2004.
3. Yusupov G.A. Modification of electropunctural diagnostics and correction of systemic pathology by wave characteristics of nosodes and homeopathic remedies. Abstract for the degree of candidate of medical sciences, 2004.
4. Yusupov G.A. Energy information medicine. Homeopathy, homotoxicology, electropuncture according to R. Voll. - M.: "Moscow news", 2000.
5. Yusupov G.A. Method for the treatment of duodenal ulcer and medicinal product for its implementation. RF patent No. 2182486.
6. Voll R. Anwendung der Medikamententestung und der Nosodentherapie zur Mesenchimentsclackung bzw. Mesenchimreaktivierung. In: Medikamententestung, Nosodentherapie und Mesenchimreaktivierung, MLV mbH-Uelzen, 1976.
7. Voll, R. Nosodenanwendung in Diagnostic und Therapie. MNV-Uelzen, 1977.

---

Yusupov, G.A. Possibilities of the modified method of electropunctural diagnostics in determining the mutual compatibility of homeopathic medicines. Yusupov // Traditional Medicine. - 2006. - No. 1 (6). - S.27-30.

[To favorites](#)