

"Peaceful atom in every home"
A. Matison
(Firma Matisons. Ltd, Riga, Latvia)

Unfortunately, this slogan, announced by the Communist Party, was really implemented, but not in the form in which it was assumed by the ideologists, but in the form of radiation contamination, which was most clearly manifested after the tragedy at the Chernobyl nuclear power plant. The most dangerous thing about this weak lesion is that it can change the configuration of DNA and cause changes in chromosomes. If massive radiation has occurred, then acute radiation sickness develops, which quickly leads to the manifestation of symptoms of acute radiation sickness, which are described in the available medical literature. But, along with massive radiation damage, there are more and more cases of weak radiation damage, which can be hidden under the guise of various progressive chronic diseases that are almost not treatable as classical methods of treatment.

Modern classical medicine has practically no methods for diagnosing radiation damage, and the analysis of determining abnormalities in chromosomes and DNA is rarely done.

VRT (vegetative resonance test), and especially VRT +, makes it possible to identify not only the radiation damage caused by the agent, but also, when measured at level 4, to determine which chromosomes are affected by this agent.

If we consider homeopathy as one of the main parts of MRI, then its application lies in the correct choice of the drug. The classical method in choosing a remedy is the repertorization of the patient, but, unfortunately, the methods of repertorization of classical homeopathy do not quite work in case of radiation injury.

The ART + method allows not only to determine the level of damage - DNA, chromosomes, but also to determine the isotope that caused this damage. To do this, you can go in two ways:

1) Use electromagnetic induction on the frequencies of the elements periodic tables through the UMT "loop", setting the amplitude to 100 units. and through the filter "primary defeat" find the primary affected element. Then, already knowing this element, find that homeopathic preparation (often from the list of radionuclides) that neutralizes this lesion and, having included it in the chain of exposure, conduct MRI according to the chosen strategy.

2) Entering the section of the drug selector - "Radionuclides", go over everything in order and find the one that will neutralize this defeat as much as possible (filter - primary defeat).

It is better to find the element and isotopes at the 3rd level of measurement of ART +, because this does not create an additional load on the cell nucleus, but the search for the induction program of BRT through the found drug at the 4th level, because our main influence in this case is aimed at correcting DNA and chromosomes.

In this case, the use of the potentiated drug will be carried out not according to the homeopathic likeness, but according to the principle of isopathy - to the agent that caused the lesion.

Before starting treatment and prescribing a drug found in this way, you need to check it for efficacy and tolerability at 2 and 3 levels of ART + measurement.

In this case, doctors who use BRT and have ART + in their arsenal have undeniable advantages in diagnosing and choosing the right effect on the patient.

Unfortunately, this method has to be used more and more often, as the damage from radioactive isotopes is expanding.

A. Matison "Peaceful atom in every home" //

"IMEDIS", 2009, v.1 - p.122-124