The BRT algorithm when exposed to the patient's body on various

levels

V. I. Reshetilovone, M.V. Vlasenkoone, M.S. Sorokopud2 (onecity of Budapest, Hungary; 2Zaporozhye, Ukraine)

The issues of adequate and most effective impact on the patient's body by the BRT method remain important for all doctors using this method.

Purpose of the study

1. Create an adequate and effective algorithm for BRT in treating patients using the effects of BRT on their body at various levels.

2. Carry out the selection of drugs, preparation of prescriptions and therapy regimens in relation to the proposed 3-level treatment regimen.

Materials and research methods

The work was carried out on APK "IMEDIS-EXPERT", software version 5.30. Medicines are recorded on homeopathic crumbs (sugar crumbs).

This algorithm for treating patients at different levels of exposure was developed in the process of practical work for three years in an outpatient setting.

Therapy according to the proposed scheme of exposure at three levels was carried out in dynamics in 500 patients with various pathologies.

Results

To carry out the most effective, adequate treatment by the method BRT

the whole body is conditionally divided according to the direction of the effect of therapy and drugs into three levels:

Level I - blood, lymph, interstitial fluid.

II level - intracellular formations.

III level - DNA.

Naturally, this division is very simplified. But it allows a professional doctor or researcher to accurately localize the direction of the therapeutic effect, and the patient is given a clear and complete understanding of what is happening to him, what drugs and why he is taking, at what level they act, what reactions from the body are possible during therapy. The proposed algorithm provides an optimum understanding of the scheme and tactics.

BRT for both the doctor and the patient. This approach has proven to be very effective and confirmed the importance of understanding what is happening from the outside. the patient.

BRT algorithm at all levelsaction on the patient's body on various

Selection	drugs,	drawing up	recipes	and schemes	therapy
-----------	--------	------------	---------	-------------	---------

in relation to a 3-tier treatment regimen.

The device is switched on as standard. We open the window "Lists of medicines". In this window, open the folder "Complexons of various companies", in it open the folder "Dr. Schimmel's preparations". Through the section of this folder "Minimum test kit" we make a test analysis of the body, first of all, paying attention to the oncological predisposition.

Level I

The main task of therapy at this level is drainage and stabilization of the state of the body at the level: blood, lymph, interstitial fluid, the outer membrane of the cell.

1.1. In the simplest version, as well as for beginners, a recipe is prepared from the section "Resoplex" (No. 1-12, 18-25), HS-complexes (No. 4-8). In addition, Bioenergetic Protection (FlowerPlex No. 5 by GUNA) must be added to the recipe. It acts as a "recipe limiter".

1.2. With enough practical skills, the recipe can be strengthened drainage preparations of the company "OHOM" (No. 62–76). Mesenchymal metabolic drainage (# 77) should not be used at the first stage of treatment.

1.3. In case of inflammation of the respiratory tract, it is advisable to add to the recipe Dr. Rekeveg preparations: 50: P49 (Preparation of the sinuses and nose) and / or 46: P45 (Preparation of the larynx).

As a result of an individually selected combination for a specific patient, recipe No. 2 is obtained, against the background of which one or another therapy is carried out, and then the therapy is assessed by the method of segmental diagnostics.

Preparation of prescription drugs # 2.

We turn on the "Bioresonance therapy" window, go to the "Drug mode test", turn on the "Select all meridians". Recording produced within 200 s, as a result, a drug is created with the selected homeopathic complexones and all 20 meridians.

The standard dosage of recipe number 2: 3 times 6 balls a day or 2 times 8-10 balls. The duration of the use of these drugs is from 2 to 4 weeks. Most often - 4 weeks.

Recipe No. 1 is technologically prepared last, although the selection of drugs is carried out much earlier. Open the folder "Meridian drugs ", section" Combined MP ". The latter are represented by three drugs: for substance intoxication; for acquired toxic information; for congenital toxic information. We denote them accordingly: Rp.1-1, Rp.1-2, Rp.1-3.

Rp.1-1 is suitable for work at the first level, it is prepared using the same technology as recipe # 2. Dosage: 1 time a day in the morning on an empty stomach, 10 balls. The course of therapy is from 2 to 4 weeks.

II level

Intracellular drainage and restoration of normal cell function. 2.1. We prepare recipe No. 2 from the "Meridian drugs" portfolio, sections "FM-complexes" No. 1-13, 15, 19-21), "Other meridian preparations (No. 1-4, 8, 13-15) and" Meridian complex preparations "(No. 1, 6, 7) ...

We prepare recipe No. 1 on the basis of "Combined MPs" - for the acquired toxic information - and denote it as recipe No. 1-2.

The technology of preparation of preparations of recipes No. 1-2 and No. 2, dosage and duration of their administration remain the same as in the first level of exposure.

2.2. If after the treatment the problems remain at this level, then do not rush to the next level. It makes sense to work horizontally.

For this, prescription No. 2 is prepared either from drugs "MP for the treatment of organs" (No. 1-9) or preparations "OHOM" ("Alfa-Omega"), group SIN (No. 1-61). Use the latter with caution, as they contain nosodes and therefore can cause exacerbation.

2.3. A completely new way of working at this level can be obtained, working with "GUNA" preparations. We work through the "Repertory" window, "GUNA" repertory, 2007. Here, for many nosologies, ready-made recipe schemes are given; the work on the selection of the chelator has already been completed.

It should be emphasized that in these recipes, the drugs are selected not only for compatibility, but also for the nature of the effect: a specific agent, a drainage agent, and PNEI (psycho-neuro-, endocrine, immunointegrative) drugs.

GUNA preparations are the so-called "avant-garde" homeopathy, as they include not only traditional homeopathy, but also genetic engineering products, as well as stem cells.

The experience of working during the year with this group of drugs, which combines natural and wave forms, has shown their very high efficiency.

The principle of work, when natural homeopathy is combined with wave homeopathy, is more physiological than the combination of wave homeopathy with biologically active additives (BAA). This is not only a theoretical point of view, it is supported by our practice.

III level

Impact on DNA

This level of work is not realistic for all patients. It is suitable for those who have sufficiently large reserves of adaptation, and this treatment will not cause excessive depletion of the body systems. First of all, we are talking about the immune system. This approach is applicable for somatic diseases, and for psychoemotional problems there are practically no restrictions. Therapy at this level cannot be carried out against the background of acute conditions.

3.1.Creation of an organism-specific nosode in the "lung" or more "Heavy" options.

3.1.1. "Easy option".

We create a nosode as follows: in the first container of the apparatus we put a glass with grains and for 440 s we carry out an organotropic one,

consistent, circular, with connecting frontal electrodes bioresonance therapy for all meridians.

3.1.2. "Hard option."

We create a nosode as follows: put a cup with crumbs in the first container and for 900 s in the "swing" mode, a fixed frequency code, a change in leads, diagonal, with the connection of the frontal electrodes, we carry out bioresonance therapy.

According to the first or second options, we get an organism-specific nosode.

This nosode is specific to this organism, but we do not clearly understand its composition.

The drug is prescribed for 1–3 days (rarely 7 days) until a standard homeopathic reaction appears. Dosage: 3 times 6 balls or 2 times 8-10 balls.

3.2. Prescription of medicinal preparations of "ONOM" company №№ 116-126.

The recording of these drugs can be carried out in three versions according to the principle: "from easy to difficult".

3.2.1. In the transfer mode, one or another drug is simply recorded in for 120 sec.

3.2.2. In the drug testing mode for all meridians prepare the selected miasmatic preparation for 200 s.

3.2.3. The drug is prepared in the drug testing mode for all the meridians with the connection of the patient through the frontal electrodes, that is, the disharmonious oscillations of the patient are also recorded. The preparation time is 200 s.

The duration of treatment and dosage vary quite a lot - from 1 time per month for 5–20 balls to 1 time per week for 5 balls - and depend on the results of individual testing.

3.3. Preparations in LM potencies and M potencies (folder "Other drugs ") can also be used in the treatment of miasms. However, they require certain skills in work and their use is not recommended for novice doctors.

Conclusions:

1. The proposed algorithm and a 3-level system of approach to treatment process in BRT have established themselves as versatile and effective

working for various forms of disease.

2. The proposed technique allows the doctor easier to navigate in everything variety of drugs presented in the selector, build individual treatment regimen, to avoid sharp exacerbations and get a good therapeutic effect.

3. With regard to the proposed 3-level BRT system to the doctor an algorithm for the selection of drugs, preparation of prescriptions and therapy regimens is provided.

4. The system of therapy at the three proposed levels has a positive effect on psycho-emotional state of patients, their contact with the attending physician and increases the effectiveness of the treatment.

V.I. Reshetilov, M.V. Vlasenko, M.S. Sorokopud BRT algorithm when exposed to the patient's body at various levels

"- M .:" IMEDIS ", 2009, vol. 2 -Pp. 123-129