

Treatment of atopic dermatitis in children

YES. Kazmirovsky, S.V. Artemenko, E.A. Igoshin, I. Yu. Shumilova, O. N.

Tkachenko, L.I. Ruchkina, N.A. Safargalina, L.V. Chumak

(Center for Bioresonance Therapy "EliteMed", Krasnodar, Russia)

Treatment of atopic dermatitis (AD) occupies a significant place in modern pediatrics, because now there are practically no children who have not had, at least once, any pathological skin manifestations.

The causes of AD include: hereditary factors, various physical, chemical and biological factors (dysbiosis, infectious toxic agents, insufficient resistance to stress, etc.).

In the modern standard treatment regimen, along with a hypoallergenic diet, corticosteroid drugs and antibiotics are included, which, as a rule, lead to a perversion of the immune response and the appearance of more serious diseases.

When examining patients with AD, we used:

1. Electropuncture diagnostics by the method of R. Voll.

2. Vegetative resonance test "IMEDIS-TEST", during which the role of various loads was clarified: geopathogenic, radioactive, psychovegetative, as well - microbiological burdens. The states of microelectrolyte and acid-base balance, enzymatic systems. Identified target organs and energy meridians with maximum disorders.

3. As an additional control, used laboratory, instrumental and bacteriological examination methods.

In the selection of a complex drug for the treatment of blood sought to pressure, we normalize BAP, leading to a rational energy and immune distribution, which ultimately affects not only skin manifestations but also the entire regulatory system of the body as a whole.

As one of the main filters, KMH was used, as it most fully reflects the energy-informational state of the patient, through which indicators of possible causes of the disease were selected.

The sum of these pointers + KMX was placed in a passive electrode and used to select medications for treatment.

An autonosode was created from the sites of pathological skin manifestations. A constitutional preparation was selected.

The selected preparations were finally potentiated through CMH before recording on the media.

A special place is occupied by drugs and induction programs to relieve psycho-vegetative loads. Also, drainages and allopathic drugs were used to eliminate dysbiosis, a hypoallergenic diet.

Clinical examples

1. Patient M., 12 years old, suffered from AD almost from birth. Long-term treatment with various specialized specialists had no success. At the time of examination: various flaky patches and nodules on the skin in the area

distal joints of the limbs, severe itching in them, areas of hyper- and hypopigmentation.

When diagnosing on method ART, identified maximum psycho-vegetative loads, disturbances in the functioning of the liver, intestines, allergic reactions with an autoimmune component. Through the KMH, medicines were selected for treatment: energy-stabilizing and constitutional drugs, drains, autonosode (from the detached affected skin), as well as allopathy (normoflorins) to eliminate dysbiosis. After starting treatment, the skin the child was cleansed in 3 days. Follow-up is 6 months.

2. Patient K., 14 years old. AD developed as a result of transplantation hay an infectious disease (severe flu) at 9 years of age.

The manifestations of blood pressure intensified at the breaking diets, slightest insignificant stress.

Large doses of corticosteroids were used to relieve cutaneous manifestations.

Diagnostics using the ART method revealed high geopathogenic and psychoemotional loads, chronic inflammation of the liver, pancreas, and general intoxication. In addition - complete blocking of 1, 3, 4, 7 chakras.

The patient underwent general detoxification, exogenous BRT (stress program, fight against a defined infection), energy stabilizing drugs (from the groups "Fateology", SDA, Aura Soma, etc.), a constitutional drug was selected, courses of endogenous BRT were carried out.

The selection and potentiation of medicines was carried out through the KMH.

The skin cleared up after 3 weeks.

Control by ART revealed a significant normalization of indicators.

Follow-up- 4 months.

conclusions

In the tactics of treating atopic dermatitis, it is necessary to take into account various etiological factors, which are specified in ART.

When selecting diagnostic and therapeutic drugs, it is advisable to take into account the general energy-informational component of each patient.

This allows prescribing more accurate and effective treatment, shortening its duration, reducing the amount of allopathic medicines taken, and optimizing the state of the body's self-regulation mechanism.

YES. Kazmirovsky, S.V. Artemenko, E.A. Igoshin, I. Yu. Shumilova, O. N. Tkachenko, L.I. Ruchkina, N.A. Safargalina, L.V. Chumak Treatment of atopic dermatitis in children // XVII