

## The use of various methods in the treatment of animals

M.A. Miller

(St. Petersburg, Russia)

It so happened that in my practice as a family doctor, pets become patients after children and their parents, grandparents. This practice began with his own dog, which in his younger years was very active, and once was injured in the forehead by a horse. Apparently, there was a concussion of a rather severe degree, since he could not shake off the water after bathing, could not open his jaw. Then I used only the homeopathic method and treated the dog with arnica 6, hypericum 6, 1 grain each, every hour for the first day. Further, more rare techniques were needed, as is customary in the case of acute conditions. On the second day, the dog returned to its previous activity.

There was still a difficult case after the deratization: housing and communal services were fighting rodents, apparently by spreading some kind of poison for the rats. Several young healthy dogs died in the house within 2 weeks. Our dog, apparently, also licked something during the walk. After that, the dog had bloody vomiting and bloody diarrhea. The classical homeopathic method also helped out. For 24 hours she gave the dog Camphor Rubini and Mercurius Corosivus. The dynamics are persistently positive by the end of the first day. Later she consulted with veterinarians about this kind of poisoning - according to their forecasts, animals die within 3 hours after poisoning with hemolysis phenomena.

In the summer of 2008, another trouble happened - a young strong boxer attacked the dog, tore his scrotum. The wound was lacerated with a skin defect and a tear in the muscles surrounding the testicles, 3-4 cm in diameter. Castration was suggested by telephone consultation with the veterinarian. Refusing the operation, she performed exogenous bioresonance therapy with fixed frequencies "trauma", "analgesic", "hemostatic". Since the dog was in serious condition, with the help of an inductor, I wrote off the oscillations ("phantom" of the animal) from it in order to conduct an examination. For a month, there was treatment with homeopathic medicines, frequency therapy. There remained a small skin defect, here preparations from the firm "Sinergia" helped the tortoise - skin and muscles that were given twice with an interval of 10 days - wound healed by primary intention.

I will cite recent observations, after which I was imbued with special respect for the methodology of A.A. Hovsepyan. At the age of 12, the dog began to show signs of malaise, depression, weakness, and most importantly, began to whine in pain when they tried to lift him. He stopped jumping into his chair at night, shortness of breath appeared.

Through the pointer to the presence of free radicals, an organ was found - the ureter, in which the amount of free radicals corresponded to BI 21, with an optimal BI 4. Using the IMEDIS equipment, I determined that it was leptospirosis. The organ-assistant was the liver in potencies D15, D30 with catabolism of 2, 3 degrees, alkaline reaction and voltage of 2, 3 degrees in the parasympathetic part of the ANS. Having put the recorded treatment chain into the load, I administered endogenous BRT to the animal for 15 minutes. She did not give any more drugs. The next day was already visible

positive dynamics, and after 2 days - there are no signs of malaise, there were no more painful reactions and shortness of breath. Leptospirosis is not tested.

The last observation was January 2011. The dog is 14.5 years old. The hind legs began to weaken.

I decided to regulate the acid-base state of the blood. Through the "Blood" index, 5 degrees of alkaline reaction were determined, placed all these degrees in inversion, determined the number of acidic valencies - the acidity of the 1st degree is tested. Through BI, I determined the initial and the optimal level of acidity in the blood. The chain looked like this:

Blood of Somr. + Alkalinity 1 degree + Alkalinity 2 degrees + Alkalinity 3 degrees + Alkalinity 4 degrees + Alkalinity 5 degrees.

Blood of Somr. + [Grade 1 Alkalinity] + [Grade 2 Alkalinity] + [Grade 3 Alkalinity] + [Grade 4 Alkalinity] + [Grade 5 Alkalinity] + Grade 1 Acidity + Mesenchyme Activity (BI) 4.

Blood of Somr. + [Grade 1 Alkalinity] + [Grade 2 Alkalinity] + [Grade 3 Alkalinity] + [Grade 4 Alkalinity] + [Grade 5 Alkalinity] + Grade 1 Acidity + Suboptimal BI + Mesenchyme Activity (BI) 11.

The modeled chain was as follows:

Blood of Somr. + [Grade 1 Alkalinity] + [Grade 2 Alkalinity] + [Grade 3 Alkalinity] + [Grade 4 Alkalinity] + [Grade 5 Alkalinity] + Grade 1 Acidity + Mesenchyme Activity (BI) 11.

I recorded this chain on crumbs, placed it in a load, the first organ-assistant determined the common bile duct with the following parameters:

Common bile duct D3 + Common bile duct D4 + Common bile duct D5 + Common bile duct D6 + Anabolic processes 4 tbsp. + Anabolic percent 5 tbsp. + Anabolic processes 6 tbsp. + Grade 3 Alkalinity + Grade 4 Alkalinity + Grade 5 Alkalinity + Grade 6 Alkalinity + Grade 1 VNS Voltage + Grade 1 VNS Voltage + Grade 3 VNS Voltage + Grade 3 VNS Voltage + Grade 4 VNS Voltage + Sympathicus D3 + Sympathicus D4 + Sympathicus D5 + Sympathicus D6 + Cellular immunity + Moderate. Art. exhausted. immune system + Stress. endocrine system. 1 tbsp. + Voltage endocrine system. 2 tbsp. + Gastrin Somr. + Glucagon Somr. + Parathyroid hormone D6 + Prostaglandon PG1 Somr.

The pancreas came to the aid of the common bile duct in potencies D10 - D30. Thus, the treatment chain looked like this:

Pancreas D10 + Pancreas D12 + Pancreas D15 + Pancreas D30 + Anabolic percent 1 tbsp. + Anabolic percent 2 tbsp. + Anabolic percent 3 tbsp. + Grade 1 alkalinity + Grade 2 alkalinity + Grade 3 alkalinity + Grade 1 ANS voltage + Grade 2 ANS voltage + Grade 3 ANS voltage + D3 vagus nerve + D4 vagus nerve + D5 vagus nerve + D3 common bile duct

+ Common bile duct D4 + Common bile duct D5 + Common bile duct D6.

BRT conducted in automatic mode along all meridians with load

the received treatment chain for 20 minutes. It is important that the animal itself determines the time of the therapy. That is, it calmly lies under the inductor until a resonance is obtained, as soon as the necessary vibrations are generated in it, the dog gets up and moves away from the device, removing the inductor from itself. The positive dynamics again clearly appeared the next day after the therapy.

There was another four-legged patient - a cat Eduard, 6 years old. The owners complained of constant urine leakage. The cat suffered from urolithiasis. When testing the Bladder D15, D30 with catabolism of 4, 5 degrees, acidity of 5, 6 degrees, VNS voltage 1, 2 degrees in the sympathetic part. In addition, blockade of adaptation reserves of 1 and 2 degrees, impairment in potassium and sodium, and tension of the endocrine system of 1 and 2 degrees in terms of renin were tested on the bladder. After modeling the chain, the cat was selected homeopathic remedies Natrium muriaticum 50, Stigmata maidis D30, Kalium iodatum 3 for 3 weeks. Three weeks later, the owners conveyed gratitude from the cat.

---

M.A. Melnik Using various methods in the treatment of animals // XVII