

Actinides - drugs of the crisis in ART

O.A. Postnikova, E.V. Postnikova

(Department of Homeopathy and Electro-acupuncture Medicine FGOU IPK
FMBA MH of Russia, JSC "INTERMED", Moscow Homeopathic Center, Moscow,
Russia)

The themes of development in the mineral kingdom are reflected in the change in chemical elements according to the periodic table of D.I. Mendeleev, according to the theory of Jan Scholten and Rajan Shankaran.

However, the matter of medicine of heavy radioactive elements has not been sufficiently developed, the topics actinides only casually reviewed by J. Scholten. According to him, actinides are drugs of a pre-agonal state and agony, when it comes to the real end of life, drugs used on the verge of life and death.

According to the authors, according to ART, these drugs in testing appear periodically, in connection with the processes of life reassessment, at moments of abrupt maturation, and signal a sharp leap in development, excessive withdrawal from life or an increase in age-related changes of a destructive nature.

According to our data, in the study by the ART method, actinides indicate an existential crisis when the question of the meaning of life and death is being resolved, when the solution of the question approaches the meaninglessness of life and the possibility of leaving it, detachment from life is visible, the tested element becomes more and more difficult, fits the more difficult the situation is, the closer to the last line. In such cases, suicide is possible and, when the most severe elements are tested, the condition can be defined as depressive. This condition corresponds to serious health problems, in particular, according to ART data, oncological diseases are often tested, when a psychological program for self-destruction, unwillingness to live, is turned on. This is a decision that is not always conscious, and maybe this is a decision of only one part of the personality. This exit is facilitated by the partial disintegration of the personality,

Some people only periodically and briefly plunge into such pessimistic detached philosophical reflections, but not linger in this one state, and then, upon exiting this state, the body is able to restore the balance of health. However, people who are prone to getting sick may have trace reactions and memory of the experience for a long time, and they will periodically plunge into these sensations again and again, then at these moments the continuation of the program of self-destruction and illness is possible.

Actinides also reflect the search for the meaning of life in adolescence, with too rapid growth or a rapid transition to a separate adult existence, as well as in other periods of crisis. With a favorable exit from the crisis, changes after the crisis lead to the formation of a renewed personality.

Such a crisis can cause various changes in life: separation from the mother, family, a sharp decrease in the level of support in the family, the sudden loss of a partner, parting with grown-up children, joining society, going to school, college, to work, change of team, job, deeply felt

achievements and failures in work and art, professional skill or achievement and loss of power, experienced negatively with the denial of the possibility of a good outcome of the situation, when, ultimately, there may be an unwillingness to live.

The crisis forces us to make a choice of a further path: self-destruction, as a denial of the previous way of life, a return to the previous point of the path, with an unresolved development task - regression, or life as the development of a new way of existence, setting new tasks and goals of life.

A characteristic feature of actinides is the comprehension of life, a view of the world from a distant point, "almost from Space", a neutral or rejecting position in relation to life, non-participation in the life process.

The gradual development of age-related changes is reflected in the change in the leading drugs towards an increase in their molecular weight, the appearance of common themes and general problems associated with age, accompanied by a decrease in the functions of various organs and systems.

Age-related changes have common features in treatment regimens, in contrast to the problems of a younger age, where the main thing will be the development of hereditary limitations and their compensation or overcoming the stops of the problems of development and maturation, formation. Problems that develop with age are quite common for many, associated with a decrease in the functions of the excretory systems, with gradually developing endocrine insufficiency, primarily, a decrease in the function of the thyroid gland and gonads. There is an increase in the level of intoxication, deposition of homotoxins in tissues, both in the intercellular space and in cells.

The tasks that are set at this stage of life are adaptation to new limited living conditions, with a decrease in physical capabilities and activity, and a decrease in working capacity. When there is a narrowing of the circle of interests, it crushes the burden of disappointments, causing depression, causing a decrease in the level of hopes and faith in the future, a stop in the movement of development, when it is necessary to set new development tasks, because the old ones are already coming to an end. In such a situation, it is important to increase the activity of organs, to cleanse the accumulated homotoxins, from radioactive heavy elements in the body. In this situation, there is a gradual transition to increasingly heavy chemical elements with radioactive decay, reflecting the state of decay in the human body and psyche. And the appointment of this heaviest element tested by ART can help bring it back to life.

Actinides can be used as a diagnostic test-indicator of ART, indicating actual existential problems, a crisis or depression, with a possible severe course of the disease, for example, an increase in the likelihood of an oncological process, or, probably, a radioactive toxic load. When testing for ART, in order to check the problems in this group, you can start testing with curium - the drug of 10 columns, which, according to the authors, is associated with equilibrium and its disturbance in a period, a row.

If it has a loading effect: Curium ↓ + drug
of the Actinoid group ↑,
then, against its background, check from the heaviest elements, starting with fermia, then

einsteinium, berkelium, then drugs after curium - americium, uranium "+" another drug of the Actinide group (1), and this drug should be prescribed to the patient or included in the BRT.

It is important not to miss it, to check the most difficult element by ART, then the treatment process will go towards alleviating the symptoms.

An interesting observation by the authors is that if a patient receives light elements of the periodic table, often a heavy element, like a roof, may be needed to balance the processes in his body, limiting progress to the lower left edge of the periodic table, where the life path ends. A heavy element here is a drug that returns the patient to the middle of the table, where normal active activity in society, achievements in human society are possible.

Light elements such as hydrogen, helium, lithium and others are developmental, conducive to maturation, growth, are used for problems of the beginning of life or developmental stops, when an adult has problems with his child's part.

Regardless of age, heavy elements indicate the end of the life cycle in relation to some area of activity. The most difficult ones, tested by ART, return a person's interest in life, as well as increase his physical capabilities, lost in the previous period, help to set new life tasks and help prepare a healthy body for this.

These tested drugs can be used in BRT as therapeutic agents, as well as diagnostic drugs in the ART method that determine the goals of therapy.

In the same way as the selection of actinides with a therapeutic effect, it is possible to test the leading drug in each row (period), optimizing the solution of the problems of each stage of life.

When choosing a therapeutic drug through ART, all the central drugs of the table (10th column) can be used for BRT and ART, separately or together as a diagnostic filter task for the body, to stabilize mineral metabolism and identify unsolved problems, problems of the mineral level, both on the physical and on the psychological plane.

Conclusion: according to the authors, it is necessary to expand the indications for the use of actinides and other heavy and radioactive elements, in addition to radioactive contamination, age problems, cancer diseases or addiction to them, depression, crisis situations.

O.A. Postnikova, E.V. Postnikova Actinides - drugs for the crisis in ART // XVIII