Influence of sun activity E. Scott-Morley (Poole, UK)

When working with patients, I pay considerable attention to identifying the body burdens with geopathogenic and electromagnetic loads. Sleep disturbance symptoms and mood swings may indicate the effect of these stresses on the patient.

When testing by the method of R. Voll, the presence of loads is manifested in high values at the end points, while high values are also recorded at the KTI. When testing by the ART method, appropriate test pointers are used. I additionally check the polarization of the biological media of the body and begin therapy with its correction. Then I examine the patient.

The sun?

I usually find a geopathogenic load in about 5% of patients. Since February 2011, the frequency of occurrence of geopathogenic load among my patients has increased to approximately 85%. High values at the end points and CTE, a positive test for geopathogenic load during ART, a violation of blood polarization in each such patient. I was looking for blood polarization test therapy. Typically, each patient has a different therapy regimen, but more than 100 patients were found to be matched with a MOPA regimen with a bandpass filter of 945 Hz to 1.25 kHz with gains of 1 to 3.

This is a rather strange phenomenon, since my patients come from different places and even from abroad, and any local influence on patients is excluded. By the end of February, I realized that the state of patients I was detecting was associated with an increase in solar activity, which is cyclical. character, and in 2011 another 11-year period began. As observers note, the modern cycle is the most intense over the past 150 years. The intensity of solar flares is so great that governments such as the UK are considering various options for protecting ground-based power grids. It is not known exactly how such solar activity will affect the ozone layer, the Earth's magnetic field. And it is likely that solar activity could be the source of a significant surge in geopathogenic load.

I observed disturbances in blood polarization in most patients from February to July 2011, when disturbances in blood polarization ceased to be detected and disturbances in urine polarization were tested en masse. Electromagnetic loading was clearly tested in patients. The mode of MORA therapy for such patients was changed to the range from 1.7 to 1.75 kHz, as well as 15–20 Hz. Is this still due to solar activity?





Fig 1. EAF portrait of geopathogenic load


Clinical example

The patient at the initial appointment, February, 2011. The general picture of the measurement results is noteworthy (Fig. 1). All results are above the normal range. A number of points marked the "fall of the arrow". All this may indicate the possibility of the influence of geopathogenic or electromagnetic loads. Revealed geopathogenic load, violation of blood polarity.





August, 2011. After the correction of the geopathogenic load, it was not revealed. During this examination, the patient had a lot of points with high values and the "arrow falling" effect (Fig. 2). Electromagnetic load is tested. Revealed a change in the polarization of urine.



Rice. 3.September 2011 The patient's condition is compensated.

When discussing solar activity, one of mythe interlocutors suggested the possibilitythe influence of HAARP on the ones I observedphenomena. I certainlyabout HAARP, butdid not consider the possibility of influencingknew this system.knew this system.

My observations, based on the results electropuncture diagnostics (EAF, ART), suggested and that change polarization of urine should correlate with the patient's response to anthropogenic electromagnetic fields. I usually associated this with the usual influence of the environment, but there were also foreigners among my patients. Perhaps it is HAARP or similar impact?

I understand that an essential part of this publication is speculation. No credible evidence of the influence of the sun or anthropogenic electromagnetic fields. However, the Aqua solis and HAARP Belastung markers were tested in most of the patients. There is no means to protect or shield such influences. Therapy for polarization disorders of biological media of the body gave a positive result, patients have a positive trend.

And the best assessment of the results of therapy is the evaluation of results therapy by the patient.

E. Scott-Morley Influence of Sun Activity // XVIII

"IMEDIS", 2012, v.2 - C.305-309