

Vaccine compensation - improves the result of bioresonance
therapy

A.F. Dolgushin, F.N. Dolgushin

(Medical Center for Innovative Technologies "Medax", Tyumen, Russia)

Using vegetative resonance test and various options bioresonance therapy on the devices of the Center "IMEDIS" for 10 years, we have developed a certain algorithm for testing and treatment, which is constantly being improved to obtain more effective medical and economic results.

The main option for vegetative resonance test (ART) and bioresonance therapy (BRT), with a predominantly endogenous technique, is carried out according to Professor A.A. Hovsepian. We consider it physiologically reasonable and comprehensible in a physician's way. A comprehensive assessment of the problem organ includes, first of all: the state of metabolic processes, in conjunction with the main integrative systems of the body, information blocks, the presence of external, internal and hereditary factors, etc. As a result of this approach, the doctor receives "super-individual" and complex information about the nature and the causal relationship of the disease of the applied patient. When the body is burdened by external influences in the form of a viral, bacterial, microcotic or parasitic, the main principle of building a normalizing physiological chain of a bioresonance drug is aimed at normalizing metabolic processes (creating "uncomfortable conditions" "for the existence of microorganisms), which makes it possible to rarely use exogenous bioresonance therapy. This minimizes the imposition of an electromagnetic load on the patient and makes it possible to obtain a more effective therapeutic result.

Since November 2014, we have additionally used the method of D.D. Tikhomirov in the form of bioresonance cleansing of the body, determination and elimination of a wider range of external loads, leading to a violation of the acid-base balance towards a decrease in RN (based on the seminar of D.D. Tikhomirov, October 2014). These measures allow the patient to quickly normalize the acid-base balance and reach a higher level of adaptation reserves. The patient will do the rest himself, taking bioresonance preparations created according to the method of Professor A.A. Hovsepian.

In our message, we would like to share the first results of the treatment of frequently ill patients who have been diagnosed with vaccine burden during ART. Vaccine prophylaxis was performed at an early and later age. The total number of patients - 16, age: from 3 to 5 years - 4 patients; 5-14 years old - 5; over 14 years old - 3 patients; adults - 4 patients.

What vaccines represent the burden:

- Viral hepatitis B - 1 (confirmed clinically and laboratory biochemical liver tests);
- Rubella - one patient;
- Poliomyelitis - 2 (one adult);

- Tick-borne encephalitis - 3 people;
 - Against tuberculosis - 9 patients.
- Allergic diseases were the reason for referral in 12 patients.

Clinical example

Elisey P., 6 years old. Date of treatment 02/18/2015 Complaints when applying for nasal congestion, dry cough. Ill since 2 years of age. Observed and treated by a pediatrician and allergist for bronchial asthma of infectious and allergic genesis. Takes symbicorting twice a day (2 doses in the morning and one dose in the evening). According to ART data performed on the hardware and software complex "IMEDIS-EXPERT", a burden of anti-tuberculosis vaccine was revealed (vaccination against tuberculosis, tuberculin D100, D200); violation of acid-base balance (acidity 1-4 degree) caused by external factors. Through the Intox 1 pointer, the following are tested: chemical burden, D30 viruses (frequencies of the Epstein-Barr D5-D30 virus, parasitic burden (frequencies of intestinal lamblia in D5-D100 potencies).

At the first stage of therapy, the selection of exogenous detoxification frequency and chemical substance for bioresonance cleansing of the body was performed.

Second stage: the procedure for eliminating vaccine burden. (Vaccination against tuberculosis + tuberculin D100, tuberculin D200) + Selected meridians. A session of endogenous bioresonance therapy was performed with the recording of drugs in the first container. Selection of the amount and frequency of taking the drug.

The third stage: recording the drug "anti-lamblia". The number of globules of the bioresonance preparation corresponding to the selected daily dosage was placed in a "loading" container.

The fourth stage: work with the selected organopreparation "Small intestine". Through its frequency spectrum, a physiological chain was tested and built with the following parameters: catabolism 1, 2 tbsp. activity + 1 degree alkalinity + VNS voltage 1, 2 degrees + sympathicus D4, D5 + endocrine system tension 1, 2 degrees + melatonin, ACTH. Indication of the tension of the immune system + low degree of tension of the immune system + cellular immunity (connective tissue deficiency 1, 2 tbsp.) + Thiamine D30 + calcium D30, silicon D30. Lymphatic congestion 1, 2, 3 degrees (Blockade of the 3rd layer) + Mental stress of the 3rd degree. Very high level 3 adaptation reserves. A session of endogenous bioresonance therapy was performed through selected meridians (Lymph, Lungs, Nervous degeneration, Allergy, Small intestine, Liver, Kidneys,

Antiviral drug No. 4.

Bioresonance cleansing on the MINI-EXPERT-T apparatus, selected by the E14 program, through a 4% soda solution using hand electrodes.

Control reception with correction of treatment after 7 days. The dynamics are positive. The state of health has improved. The cough has stopped. The inhaler is used once a day (starting from the third day of treatment), moderate nasal congestion remains.

According to ART data: acidity of the 1st degree, vaccine burden is not

determined, not tested for Epstein-Barr virus frequencies. An imbalance of the immune system with an autoimmune component is determined on the nasal mucosa. Endocrine system tension 3 tbsp. hormones: adrenaline, histamine, antidiuretic hormone. There is a burden of intestinal lamblia. The pathological chain was corrected, a session of endogenous BR-therapy was performed, with the recording of bioresonance drugs. The drug No. 1 (elimination of vaccination burden) was canceled. After 7 days, according to my mother, the condition is good. Nasal breathing is free, there is no cough. Does not use an inhaler.

conclusions

Vaccinal aggravations are common. It is advisable to include a test for vaccination burden in the algorithm of the autonomic resonance test in patients of all age categories.

Elimination of vaccine burden significantly improves the result of bioresonance therapy carried out on the apparatus of the Center "IMEDIS".

Dolgushina, A.F. Compensation for the burden of the vaccine - improves the result of bioresonance therapy / A.F. Dolgushin, F.N. Dolgushin // XXI International Conference "Theoretical and Clinical Aspects of the Application of Bioresonance and Multiresonance Therapy". - M.: IMEDIS, 2015. -- P.380-383.

[To favorites](#)