

Autoallergies and behavior of children  
A.N. Mathison  
(Firma "Matisons" SIA, Riga, Latvia)

Every year, the number of children with behavior that differs from the norm is increasing, which can manifest itself as a syndrome of hyperactivity, inattention, various obsessive movements, neuroses and neurasthenia, a decrease in the assimilation of educational material at school. Often such children are recorded as inveterate hooligans, trying to influence them with pedagogical and not very pedagogical methods, which most often do not bring the desired result. The "label" hung on such a child begins to persecute him as a "sentence" already from kindergarten and goes to school, even from the side of the closest people - his parents. Well, if this child is diagnosed with any disease, then his behavior can be attributed to this disease or a side effect of treatment.

It is much worse when such a child has no visible diseases and his child is referred to a psychologist. Often, the psychologist also fails to get the desired result and to identify the cause of his inappropriate behavior, and then he is sent for consultation to a psychiatrist who tries to correct his behavior with the help of sufficiently powerful medications. At the same time, the child continues to experience quite strong pressure from the society, misunderstanding in the family. In fact, he would like to behave as required of him, but the constant irritation of H3 histamine receptors, 40% of which are in the brain, caused by allergies, and more often by an autoallergic process, does not allow him to behave differently, which causes so many complaints from the people around him.

The main mediator of allergy is histamine, which is produced by mast cells and acts on cells that have corresponding receptors. There are three types of histamine receptors:

- H1 - Muscles, endothelium, central nervous system (postsynaptic part);
- H2 - Parietal cells;
- H3 - Central and peripheral nervous systems (presynaptic part).

Thus, any allergic reaction of the body, first of all, will affect the central nervous system, since there is the largest concentration of histamine receptors, and through the work of the brain on human behavior. At the same time, the child may have almost no symptoms of diseases that we are used to attributing to allergies: itching and rashes, runny nose with frequent sneezing, bronchospasm, rheumatism and others. In blood tests, he may not have an eosinophilic reaction, and the blood count may be normal. And with all this, we can reveal a violation in this child when testing with the help of electropuncture diagnostics according to the method of R. Voll, which is manifested by decreased or greatly increased indicators of measuring points on the meridian of allergy, bladder and lymph. And also investigate using a vegetative resonance test (ART),

more accurately allows you to diagnose problems in the body. The readings of the second level of measurement of the VRT + device will be associated with the state of the cell membrane, which most often manifests itself as a rapid allergic reaction, and when diagnosed at the third level, violations in the cytoplasm of the cell, which correspond to a delayed allergic reaction.

Even if we do not detect any abnormalities during the diagnosis using the ART method on the allergy meridian, this does not mean that the patient is not allergic. Entering the ART section "Allergy", it is better to test both food allergies (more or less sensitive) and auto-allergic processes. With a positive auto-allergy test, you need to find an organ that will experience the greatest allergic load. It is also necessary to identify all allergens that may cause an allergic reaction, and through the pointer to the primary lesion (Zin. Met. D400) find the most important (triggering) allergen.

Since allergy is a perverted immune response, both cellular and humoral immunity play an important role in it. And, of course, it is necessary to test the state of lymphocytes, thymus and spleen as the main immune organs from potency D3 to potency D30.

Since most often disturbances in the immune system occur due to vaccinations, helminthic invasion, and more recently - more often due to a radioactive load that causes changes in the immune system, then they need to be tested as well.

If, with a simple food allergy, we can use for desensitization revealed with the help of Zin. met. D400 allergen, in the correctly selected potency, having included it in the therapeutic circuit, then cases with an identified autoallergic reaction cause some difficulty.

The following treatment tactics can be proposed. If during an autoallergic reaction a radioactive load is also detected, and when testing lymphocytes, thymus and spleen, their low potencies are revealed, then using the device for magnetic therapy "loop" in EPT mode, it is necessary to test the entire periodic table and find the element corresponding to the highest indicator of radioactive load and the lowest potency of lymphocytes or thymus. After that, in the selector, select the potency of this element or its salt, which will compensate for all of the above pathological indicators. If there is no such drug in the selector, then you can test the element closest to the periodic table and start treatment in the mode of induction BRT or MRI of allergies with the inclusion of this drug in the therapeutic circuit.

If, with a positive test for auto-allergy, there is no radiation exposure, then one of the drugs: Salmon (salmon), Salmon (trout), Gadus morhua (cod), Homarus (lobster), Cenchris (moccasin snake) in the correct potency will help to cope with auto-allergies.

After the end of the session, we write on the homeopathic crumbs the drug that was used in therapy, test the optimal dose and give it at the end of the intake. Additionally, repeated doses of the drug are prescribed.

Interestingly, the behavior of patients begins to change from the first day after

therapy. And these "hooligans" become diligent and obedient, their academic performance improves and conflicts in the team and at home are sharply reduced.

Of course, a prerequisite for treatment is the exclusion from the diet of all those foods that were previously identified as allergens during testing.

Such patients should be invited for an appointment approximately once every two to three months at the beginning of treatment, and as they recover, the interval increases to six months. It should be noted that the allergic reaction is not constant, but can change over time. Therefore, every time at the reception it is necessary to carry out a complete examination.

Such a correction with the help of multiresonant BRT allows children, and not only children, to become adequate in society and to fully reveal their potential and talent.

Unfortunately, most teachers, psychologists and psychiatrists are not informed about the existing problem of autoallergy and try to solve it based on their own knowledge. Appointment of the correct diet, homeopathic remedies and bioresonance effects can radically change this situation.

---

Matson, A.N. Autoallergy and behavior of children / A.N. Matson // XXII International Conference "Theoretical and Clinical Aspects of the Application of Bioresonance and Multiresonance Therapy". - M.: IMEDIS, 2016. -- S.304-307.

[To favorites](#)