Statistical data on the application of the method of bioresonance therapy in medical practice L. B. Kosareva, S.I. Fedorenko (Center "IMEDIS", Moscow, Russia)

The generalized statistical data (Table 1) are compiled on the basis of the work of doctors at 8 medical centers during 2016. The statistical data are provided in accordance with the proposed form and are grouped by the ICD-10 disease classes.

In total, 20611 patients were treated for various nosological forms. A significant improvement was noted in 53.4% of cases; improvement - 43.4%; without improvement - in 3.2% of cases.

Form of nosology		The effectiveness of the treatment		
		Significant improvement	Improvement	Without improvements
Infectious and parasitic diseases	7100	4598	2379	123
Neoplasms	427	173	199	55
Diseases of the blood, blood-forming organs and certain disorders involving the immune mechanism	361	227	128	6
Diseases of the endocrine system, eating disorders and metabolic disorders	1704	693	901	110
Mental and Behavioral Disorders	805	446	311	48
Diseases of the nervous system	244	44	194	6
Diseases of the eye and its adnexa	434	246	185	3
Diseases of the ENT organs and mastoid	217	131	78	eight
Diseases of the circulatory system	741	287	445	nine
Respiratory diseases	1594	1031	483	80
Diseases of the digestive system	2264	971	1244	49
Diseases skin and subcutaneous fiber	1545	602	864	79
Diseases of the musculoskeletal system and connective tissue	1233	530	677	26
Diseases of the genitourinary system	1674	833	787	54
Complications of pregnancy, childbirth and the postpartum period	eleven	2	nine	0
Separate states, arising in the perinatal period	2	2	0	0
Congenital anomalies (malformations), deformities and chromosomal abnormalities	6	4	2	0
Symptoms, signs and abnormalities identified with clinical and laboratory research, not classified elsewhere	4	0	4	0
Injury, poisoning and some other consequences of exposure to external causes	245	187	54	4
TOTAL	20611	11007 53.4%	8944 43.4%	660 3.2%

A significant improvement was understood as either complete recovery, or the onset of

stable remission during a chronic disease; improvement was understood as an improvement in the general condition, the transition of the disease to a subacute state, accompanied by the presence of basic symptoms.

The main method of treatment was exogenous and endogenous bioresonance therapy, which was used as an independent method, as well as in combination with other methods of treatment.

Additional statistical information can be found in other articles of this collection.

Kosareva, L.B. Statistical data on the use of the method of bioresonance therapy in medical practice / L.B. Kosareva, S.I. Fedorenko // XXIII International Conference "Theoretical and Clinical Aspects of the Application of Bioresonance and Multiresonance Therapy". - M.: IMEDIS, 2017.-- P.98-99.

To favorites