

Clinical case
Anvarul Azim MD
(LLC "Doctor Azim", Moscow, Russia)

For many years, at the Doctor Azim LLC clinic, we have been treating patients with intractable eye diseases with a combined method of bioresonance therapy, acupuncture and color therapy, such as: optic nerve atrophy, vascular and degenerative retinal pathologies, glaucoma (primary, asymptomatic and advanced) , progressive myopia (myopia), farsightedness (hyperopia), accommodation spasm, asthenopia, amblyopia, double vision, astigmatism, strabismus, ptosis, nystagmus, "computer vision syndrome", "dry" eye syndrome, neurophthalmology, rehabilitation after tumor removal, cysts brain, etc.

Exogenous bioresonance therapy (BRT) with fixed frequencies, induction therapy with brain rhythm frequencies and endogenous BRT along the meridians (organs and systems) are carried out. Exogenous and endogenous bioresonance therapy is carried out on the apparatus "MINI-EXPERT-DT", "MINI-EXPERT-T", "IMEDIS-BRT-A" of the Center "IMEDIS" according to the method of Yu.V. Gotovsky.

When carrying out this complex of treatment, positive results were achieved in almost 100% of cases, which we reported at numerous conferences both in our country and abroad.

In this article, I would like to present one of the most striking cases from our clinical practice.

Clinical example

Patient, 77 years old. Examined by the FGKU CCGV of the Federal Security Service of the Russian Federation Ophthalmology department.

Examined in the department: Visual acuity Vis OD sph - 2.0 cyl -1.0 ax 78 = 0.2. OS 0 (zero), (light perception is). IOP - 20 mm Hg. Art. OU.

Diagnosis: OD - open-angle III A glaucoma, maculofibrosis, initial cataract, OS - terminal compensated operated glaucoma, cyst-altered filtration cushion, secondary keratopathy.

The patient was recommended to continue instillation of antihypertensive drops and consultation with a vitreoretinal surgeon in connection with macular fibrosis. There was no improvement in vision. Then the patient started looking for alternative methods of treatment and turned to my clinic for help.

On admission: Vis OD = 2.0 with correction, OS = 0. The first course of treatment (10 sessions) with a complex method of bioresonance therapy in combination with acupuncture and color therapy was carried out from 13.02 to 22.02.2017. After the first course of treatment, subjective improvement was noted vision, decreased visual fatigue. After the end of the course of treatment, there was an increase in visual acuity Vis OU = 4.0 and OS = 0.05 (peripheral vision appeared). At the end of the treatment, a significant decrease in the threshold values of the light sensitivity of the macular region of the retina was noted. Also, there was an improvement in general well-being, a decrease in irritability, and a normalization of sleep.

To consolidate and improve the achieved result, a repeated course of treatment was recommended after 6 months.

Monitoring continues.

Anvarul, Azim A case from clinical practice / Anvarul Azim // XXIII International Conference "Theoretical and Clinical Aspects of the Application of Bioresonance and Multiresonance Therapy". - M .: IMEDIS, 2017 .-- S. 121-122.

[To favorites](#)