

Use of peptides in ART and BRT

O.A. Postnikova, E.V. Postnikova

(Department of Homeopathy and Electropuncture Medicine IPK FMBA MH RF,
Moscow, Russia)

We worked with low molecular weight peptides cytomax from tissues and organs of young calves, developed by the Institute of Gerontology in St. Petersburg. We used, among other things, homeopathized potentiated preparations, which are close in action to potentiated organ preparations in low dilutions. When tested by the ART method, the synthesized peptides do not give a positive response, which may mean that they have some side effects, so we used natural peptides, which gave the best positive effect during ART.

Such homeopathic medicines have a striking and rapid clinical effect. The scheme for the use of peptides was worked out, in which each subsequent drug, its dose or potency, and the frequency of administrations were determined by testing.

The advantage of homeopathized preparations, even in low potencies, is the training nature of their effect, in contrast to material doses, when the preparation itself works directly. In a homeopathized form, these drugs increase and develop their own capabilities of organs and tissues to restore and maintain their normal functions, the body's reserves are raised, and the state of health improves. This is their main advantage, since in this case addiction cannot develop.

The effect of homeopathized peptides is harmonizing, antidegenerative, stimulating, it is similar to the antidegenerative effect of organopreparations in low potencies. However, since they are made from the organs and tissues of calves, and not the fetus, the embryo, like a number of organopreparations, their effect is milder and safer.

These drugs can be used for diagnostics in ART. During testing, a sequential program unfolds: which drug is more relevant at the first stage, then the next one, and further along the chain to the organ-source of the problem, which is the least amenable to therapeutic action. In this version, the body's need is revealed not for one or two drugs, but for strengthening all the weak points of the body in a certain sequence. The frequency of admission is usually 1-2 times a week for chronic processes. Only sometimes, with more acute and pronounced conditions, it can reach twice a day.

The surprising and rapid efficacy of these peptide preparations was noted.

The following were observed: the effect in type 2 diabetes mellitus when using a potentiated liver drug, the hypotonic effect of many drugs, the disappearance of signs of heart failure, arrhythmias, when using a heart drug, toning of the bladder sphincter with urinary incontinence,

a diuretic effect even with renal failure when using a kidney drug, a positive effect in multiple sclerosis and others. Improvement in general vitality and mood was observed. The treatment program requires the use of pineal gland and brain preparations to restore the levels of higher regulation.

In the scheme of using peptides, it is possible to alternate or combine potentiated with native preparations.

The effect of cytomax is very effective and fast, but it is desirable to carry out a more complete treatment program for patients. The action of these drugs is not complete without the use of homeopathic remedies, which restore the most subtle and precise, deep mental regulation; drainage of toxins and regulation of metabolic processes; precise immunostimulation using nosodes; a more complete effect of organ preparations in chronic processes.

When using native and potentiated peptides in bioresonance therapy (BRT) on IMEDIS devices, the effectiveness of treatment is significantly increased.

Postnikova, O.A. The use of peptides in ART and BRT / O.A. Postnikova, E.V. Postnikova // XXIII International Conference "Theoretical and Clinical Aspects of the Application of Bioresonance and Multiresonance Therapy". - M.: IMEDIS, 2017. -- S. 179-181.

[To favorites](#)