The use of a vegetative resonance test for psychocorrection of patients with oncopathology A. Alimbekova (Alma-Ata, Kazakhstan)

Most of the oncological diseases are based on a psychogenic factor, and its priority importance in the onset of the oncological process is now considered a proven fact [1, 2, 3]. In the scientific literature, there are many publications devoted to the influence of the state of mind on the development, as well as the course of cancer. Psychogenic carcinogens include, first of all, traumatic situations, long-term states of melancholy, despair and depression. The leading role in increasing the synthesis of atypical cells belongs to the loss of interest in life. There is a close connection between the development of oncopathology and constitutional-depressive personality structure, premorbid infantilism, low socio-economic status [4, 5, 6]. Additional changes in psychological status occur in the presence of cancer,

Currently, psycho-oncology is developing in three directions:

- study of the significance of the psychogenic factor in the emergence and development oncological diseases;

- mental changes in oncology;

- development of psychocorrection for this pathology.

In the early 1980s, Ryke Geerd Hamer discovered and described the following phenomenon: in acute conflict (or as a result of unethical behavior, note - ours), a large amount of energy is released from the surface of the cerebral cortex with a longterm fixation of this process. The trail of this catastrophe for the body can persist for years on the surface of the cerebral cortex and continue with the destruction of the organ. Conflict shock affects a specific, predetermined area of the brain. The effects of shock can be seen in the CT scan as a set of clearly visible concentric circles. Conflicts of the same type always affect the same area of the brain. Mr. Hamer studied in detail the phases of transition of unresolved shock and stress, discovered five biological laws, as well as the process of natural healing from disease.

In our study, we decided to rely on Hamer's conclusion: "Cancer is one of the consequences of an unresolved 'unexpected isolated conflict shock unfolding simultaneously in the psyche and the brain" (cited by Caroline Margolin). A group of specialists in Ethics of Behavior (Novosibirsk), using the method of identifying negative programs in consciousness during an ethical conversation, have achieved good results in the treatment of cancer patients. At the same time, they used the method of searching for the root cause of the problem, a conscious introspection of one's life and correcting erroneous decisions, delusions made in life by the person himself.

During the vegetative resonance test to assess the initial state of the patient, the following test indicators were used: China, C1000, False polarity, Hypothalamus D800, Epyphisis D4 1–5 cu, the indicators of the scale "Morphology" - "Morphological processes at the intracellular level" ...

We have put forward the assumption that in order to eliminate energy information programs (EIP) it is possible to narrow the field of research if we accept SHOCK and its consequences as a markercreating conditions for the moment of forcible introduction of programs with pathological

information in the information field of a person. The emphasis in the search should be placed on finding the moments of the patient committing the most serious ethical violations, which in the history of mankind were punished by death. These are the so-called "7 deadly sins".

We proposed to view this situation as a man-made ground for an attack. The work consisted of preliminary diagnostics by the ART method - identifying frequency indicators for the presence of an oncological process, EIP, various types of burdens, establishing a relationship between them and cases of severe shock, shock, with situations of violation of Ethics (agreement with erroneous decisions in life situations that harmed a person or others) ... When conducting a search for worldview errors, it is important to pay special attention to identifying externally imposed programs, blocks, causes of energy loss, which optimizes the process of correcting and correcting a pathological condition.

In the course of the analysis of the problem, the subsequent working out by the patient of the indicated problem, a repeated ART examination with a comparative analysis is carried out. If the identified problem is identified correctly and realized by the patient, voiced, then the ART indicators change. A favorable psychological resolution of an old conflict not only eliminates shock damage, but can also trigger the body's self-healing mechanism. Every patient with cancer retains a painful memory of the shock situation, the most difficult time in their life, for life. All patients were searched for a chain of events related to shock, and a detailed analysis of the causal relationship, the essence and assessment of the harm caused, and a correct reassessment of the situation was given.

Reconsideration of the psychosomatic causes of the disease - mistakes in thinking and behavior in choosing solutions to life problems - led to the elimination of testing of Chin C1000, false polarity, Deep mental processes (Hypothalamus D800), to self-healing of the state of Soul and Spirit, to the realization of the need for a meaningful transition to the rejection of violence , to the creation of goodness, love for oneself and people. It was these factors that became the healing principle. Rejection of erroneous ideological concepts, a conscious change as a result of this own character, rejection of erroneous behavior, not in words, but in practice, led to the complete disappearance of EIP testing.

Clinical example

Table 1 shows the indices of the morphological scale (Acidum fluoricum AF) obtained during the initial examination of patient M. by the ART method and the indices of the repeated examination carried out after the "ethical conversation".

Table 1

Diagnostics by morphological scales (Acidum fluoricum AF)

Acidum fluoricum AF	23.07.15	morphology	25.07 morning	25.07	25.07
					13 h
35 the oncological process is not	80/40	+	80/80	Introspection,	80/80
eliminated by the EIP program	there is			changing yourself.	
34 cancer process	80/40	+	80/80	Work on	80/80
	there is			eliminating	
33 cancer process	80/80	elimination	80/80	causes	80/80
32 oncological process in situ	80/80	NP - us-	80/80	family quarrels and	80/80
		trenching		emotions,	
31 preconf. process 2 tbsp.	80/80	mistakes	80/80	scandals,	80/80
30 preconf. process 1 tbsp.	80/80	behavior	78, yes	mistakes	80/80
29 the presence of good quality.	80/60	with the parent-	60 there	education	60
tumors		lyami,		children.	
28 presence of benign	80/80	outsiders	80/80		80/80
processes (cysts, myomas,		from 1-30 years old			
hyperplastic elements.)					
27 benign processes			80/80		80/80
(papillomas, polyps					
26 compensatory	Yes		80/80		80/80
hyperplasia					
NORM AF 25-21	80/80				80/80

Conclusions:

1. When carrying out psychocorrection of patients, the use of the ART method gives the ability for the doctor to monitor the effectiveness of the therapy.

2. Most often, in patients with oncopathology during ART a positive reaction is revealed to the following test pointers: China C1000, false polarity, Psychic load Epyphisis D4, a pointer to the presence of deep mental processes (Hypothalamus D800).

3. Changes in indicators (towards improvement) in the course of ethical conversation means the patient's awareness of the problem and the study of this block of the psyche.

Literature:

1. Simonton K., Simonton S. Psychotherapy of cancer. - St. Petersburg: Peter; 2001.

2. Kozlova N.V., Androsova T.V. Socio-psychological support

cancer patients // Bulletin of the Tomsk State University. 2010; 335: 142-7.

3. Arkhipova I.V., Kokorina N.P., Magarill Yu.A. Psychogenic factors and breast cancer glands // In the book: Actual problems of breast cancer. Kemerovo; 2003; no. VII: 7-8.

4. Marilova T.Yu. Psychological features of cancer patients // Bulletin

Russian Cancer Research Center named after N.N. Blokhin Russian Academy of Medical Sciences. 2002; 3: 47-51.

5. Gumenyuk L.N., Roslyakova V.A. Biopsychosocial risk factors for formation maladjustment in patients with malignant tumors of the maxillofacial region // Journal of Psychiatry and Medical Psychology. 2010; 1-2: 103-6.

6. Ivashkina M.G. Experience of psychocorrectional and psycho-rehabilitation support personality in conditions of oncological disease // General Medicine. 2010; 3: 49-54.

Alimbekova, A. The use of vegetative resonance test for psychocorrection of patients with oncopathology / A. Alimbekova // XXIII International conference "Theoretical and clinical aspects of the use of bioresonance and multiresonance therapy". - M .: IMEDIS, 2017 .-- S. 345-349.

<u>To favorites</u>