

Bioresonance therapy in the treatment of adenoids  
A.F. Dolgushin, F.N. Dolgushin (LLC  
"MCIT" Medax ", Tyumen, Russia)

Adenoids (adenoid growths, adenoid vegetations) are a pathological increase in the size of the pharyngeal tonsil. The pharyngeal tonsil is located in the fornix of the nasopharynx and is part of the lymphoid pharyngeal ring of N.I. Pirogov. Adenoid vegetations are caused by the proliferation of lymphoid tissue of the pharyngeal tonsils. They are widespread in children, most often observed between the ages of 3 and 7 years.

The development of adenoids is facilitated by diseases that cause inflammation of the nasal mucosa and tonsils (measles, scarlet fever, herpes viruses, etc.).

In orthodox medicine, instrumental methods are used to diagnose adenoids: X-ray, computed tomography, endoscopic methods, posterior rhinoscopy - a classic method of examination through the mouth using a special mirror, the surface of the adenoids is visible with anterior rhinoscopy (examination through the nose).

There are 3 degrees of growth of adenoid vegetation. Inflammation of the adenoids - adenoiditis can be acute or chronic. Enlarged adenoids of 2-3 degrees can lead to difficulty in nasal breathing, impaired drainage function of the Eustachian tube and inflammation of the middle ear, hearing impairment, snoring during sleep, delayed physical and mental development.

Conservative treatment used in otolaryngological practice is often symptomatic. Surgical treatment does not always cure relapse.

In our practice, patients with adenoids accounted for 70% of all children who applied. Basically, these are often and long-term ill children. Some turned to the referral of the local pediatrician.

In the selector device "IMEDIS-EXPERT" there is an organopreparation "Adenoids Comp." and a potentiated organopreparation "Adenoid vegetations".

The testing algorithm includes the classic and modernized versions of the autonomic resonance test (ART).

We test external loads from classical ART: geopathogenic, radioactive, electromagnetic. If the test is positive, we must look at what organopreparation the revealed load and adaptation reserves correspond to.

1. Testing for vaccine burden. With a positive response
  - determination of causal vaccination and its compensation.
2. The presence of mental stress. If available - selection of Programs rhythms brain, preparations "Bach Flowers"
3. Presence of mesenchyme blockade, with which organopreparation it is related; in the presence of a blockade, its elimination by creating a private bioresonance preparation.
4. Organopreparation "Adenoids som." or "Adenoid vegetations" in tested potency.

5. Determination of the available biological index and optimal BI via Cu met. D400. We leave the optimal BI in the selected preparations. Using the initial biological index, we determine the cause of slugging of the adenoids (lymphoid tissue) Intox I, Intox II, Intox III. Intox I is tested more often. Using pointers to toxic loads from "METABOLIX" preparations of the "MEDFARMA" group, we identify the existing causes of toxic loads, including those caused by bacterial, viral, mycotic or parasitic infections. We determine the nosodes of specific pathogens, most often it is the Epstein-Barr virus. Through the pointer to the organopreparation "Adenoids comp." + the available biological index, we select a program to optimize the detoxification of the body using contact electrotherapy (for children over 7 years old).

We form a pathophysiological chain according to A.A. Hovsepyan, which allows assessing the metabolism, the state of the autonomic nervous, immune and endocrine systems, deficiency / excess of trace elements and vitamins, the degree of lymphatic congestion and connective tissue insufficiency. We translate the chain into the optimal one, setting the optimal biological index at the end of the chain. We conduct a session of basic endogenous bioresonance therapy with subsequent recording of a bioresonance preparation. The organopreparation of the second chain is more often determined by the liver or pancreas.

To inactivate the detected pathogen in the presence of indications, we use the technique of multiple inversion of the pathogen nosode (Workshop by D.G. Bocharova, December 2016, CIMS "IMEDIS" LLC). The use of multiple inversion of the drug made it possible to reduce the duration of treatment and the number of procedures for exogenous bioresonance therapy. Control testing is performed for all patients in 7-10 days. As a rule, further intake of antiviral, antibacterial bioresonance drugs is inappropriate.

To improve the state of the immune system, which not only provides protection against infectious factors and immune surveillance, but plays an important role in the prevention of diseases, we include a natural immune corrector Transfer Factor from 4Life Research (USA) in the complex treatment. Including it in treatment allows you to get faster rehabilitation, protects the child from infectious diseases in the future.

#### Output

Vegetative resonance test carried out on the APK "IMEDIS-EXPERT" allows you to determine the cause-and-effect relationship in the development of the disease. Bioresonance therapy in combination with immunocorrection Transfer-factor contributes to the implementation of the ideal principle of healing: treatment - rehabilitation - prevention.

[To favorites](#)